

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44282 A2

- (51) International Patent Classification⁷: **C07K 14/00**
- (21) International Application Number: **PCT/US00/33793**
- (22) International Filing Date:
13 December 2000 (13.12.2000)
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/461,641 14 December 1999 (14.12.1999) **US**
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/461,641 (CIP)
Filed on 14 December 1999 (14.12.1999)
- (71) Applicant (for all designated States except US): **THE BURNHAM INSTITUTE [US/US]; 10901 N. Torrey Pines Road, La Jolla, CA 92037 (US).**
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **REED, John, C. [US/US]; 17044 El Camino Real, P.O. Box 137, Rancho Santa Fe, CA 92067 (US). GODZIK, Adam [PL/US]; 9184 Buckwheat Street, San Diego, CA 92129 (US).**
- (74) Agents: **CADENA, Deborah, L. et al.; Campbell & Flores LLP, 7th floor, 4370 La Jolla Village Drive, San Diego, CA 92122 (US).**
- (81) Designated States (national): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
— *Without international search report and to be republished upon receipt of that report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/44282 A2

(54) Title: **BCL-G POLYPEPTIDES, ENCODING NUCLEIC ACIDS AND METHODS OF USE**

(57) Abstract: The invention provides Bcl-G polypeptides and encoding nucleic acids. Bcl-G polypeptides include Bcl-G_L and Bcl-G_S. The invention also provides mouse Bcl-G. The invention also provides vectors containing Bcl-G nucleic acids, host cells containing such vectors, Bcl-G anti-sense nucleic acids and related compositions. The invention additionally provides Bcl-G oligonucleotides that can be used to hybridize to or amplify a Bcl-G nucleic acid. Anti-Bcl-G specific antibodies are also provided. Further provided are kits containing Bcl-G nucleic acids or Bcl-G specific antibodies. Such kits and reagents can be used to diagnose cancer, monitor response to therapy, or predict the prognosis of a cancer patient. The invention additionally provides methods of modulating apoptosis using Bcl-G polypeptides, encoding nucleic acids, or compounds that modulate the activity or expression of Bcl-G polypeptides. The methods for modulating apoptosis can be used to treat diseases such as cancer.

BEST AVAILABLE COPY

**BCL-G POLYPEPTIDES, ENCODING NUCLEIC ACIDS AND
METHODS OF USE**

This invention was made with government support under grant number GM60554 awarded by the National
5 Institutes of Health. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

The present invention relates generally to regulation of programmed cell death and more specifically
10 to molecules that promote programmed cell death.

In essentially all self-renewing tissues, a balance is struck between cell production by mitogenesis and cell loss due to programmed cell death, thereby maintaining total cell numbers within a physiologically
15 appropriate range. In pathological conditions, however, the balance in cell production and cell loss can be disrupted. In cancer, for example, an increased amount of cell production due to a shortened cell cycle time or a decreased amount of cell death due to dysregulation of
20 a programmed cell death pathway results in the growth of a tumor.

With regard to programmed cell death, a variety of stimuli, which occur either external or internal to the cell, initiate a pathway that ultimately results in
25 apoptosis of the cell. As is common for most signal transduction pathways, the various different stimuli that induce apoptosis likely initiate the process of programmed cell death through specific pathways. However, most if not all of these initial pathways
30 converge at a common point that generally involves a

member of the Bcl-2 family of proteins.

The Bcl-2 family of proteins regulate a distal step in the evolutionarily conserved pathway for programmed cell death and apoptosis, with some members of this family functioning as suppressors of cell death (anti-apoptotic proteins) and other members functioning as promoters of cell death (pro-apoptotic proteins). Overexpression of the anti-apoptotic protein, Bcl-2, for example, blocks neuronal cell death that otherwise is induced *in vitro* by various stimuli, including neurotrophic factor withdrawal, various oxidants, glucose deprivation, certain neurotrophic viruses, and amyloid β -peptide. In addition, Bcl-2 is overexpressed in some tumor cells and, in part, may contribute to tumor growth by altering the balance between cell division and cell death.

The Bcl-2 family of proteins are critical regulators of pathways involved in apoptosis, acting to either inhibit or promote cell death (Reed, Nature 387:773-776 (1997); Green and Reed, Science 281:1309-1312 (1998); Reed, Oncogene 17:3225-3236 (1998); Reed, Curr. Opin. Oncol. 11:68-75 (1999)). The Bcl-2 family members can be divided into two groups, those with anti-apoptotic activity, including Bcl-2 and Bcl-X_L, and those with pro-apoptotic activity, including Bax and Bak.

Four distinct domains have been identified in Bcl-2 family members, designated BH1 to BH4. The BH4 domain is a domain that mediates interactions with a variety of cellular proteins (Reed, *supra*, 1998). The BH1, BH2 and BH3 domains form a binding pocket for dimerization with other Bcl-2 members having a BH3 domain, which also functions as a ligand that binds to

the dimerization binding pocket. The dimerization function of the Bcl-2 members is an important mechanism for regulating apoptosis in that heterodimerization of pro-apoptotic Bcl-2 members with anti-apoptotic Bcl-2 members can regulate the cellular apoptotic pathways. Some Bcl-2 members only have a BH3 domain and therefore function as trans-dominant inhibitors of anti-apoptotic proteins such as Bcl-2 and Bcl-X_L (Reed, *supra*, 1998).

Another function of Bcl-2 members is the formation of ion channels. Bcl-2 members can localize to the mitochondrial membrane, and the formation of ion pores that alter the permeability of mitochondria is thought to be an important signaling mechanism for the induction of apoptosis. Thus, Bcl-2 members use at least three mechanisms to regulate apoptotic activity: dimerization with Bcl-2 members, formation of ion pores in mitochondria, and binding to non-Bcl-2 members that function as signaling molecules.

In comparison, overexpression of the pro-apoptotic protein, Bax, for example, promotes cell death when triggered by a variety of inducers of apoptosis, including growth factor withdrawal, ionizing radiation, and anti-Fas antibody. In addition, elevations in Bax expression occur in association with cell death induced by a variety of stimuli, including neuronal cell death that occurs due to ischemia, epilepsy, spinal cord injury, and certain neurodegenerative diseases such as Parkinson's disease and Alzheimer's disease.

Although aberrant expression of members of the Bcl-2 family of proteins is associated with various pathologic conditions, the mechanisms by which these

proteins mediate their action is not known. Often, the action of a protein can be inferred from its structural relationship to other proteins, whose functions are known. However, while the Bcl-2 family proteins share
5 certain structural homologies with each other, they do not share substantial amino acid sequence homology with other proteins, further hindering attempts to understand how the Bcl-2 family proteins such as Bcl-2 and Bax regulate cell death.

10 Thus, a need exists to identify proteins involved in the programmed cell death pathway and to identify methods of regulating programmed cell death for therapeutic applications, including treatment of cancer. The present invention satisfies this need and provides
15 related advantages as well.

SUMMARY OF THE INVENTION

In accordance with the present invention, there are provided Bcl-G polypeptides and encoding nucleic acid molecules. The invention polypeptides and encoding
20 nucleic acids are useful for modulating apoptosis by altering the expression or activity of Bcl-G. The Bcl-G polypeptides and encoding nucleic acids can be advantageously used to diagnose or treat cancer, in particular prostate, ovarian and leukemia. Furthermore,
25 the Bcl-G polypeptides and encoding nucleic acids are useful to generate or screen for agents that can alter Bcl-G activity or expression, which can further be used to treat cancer. Bcl-G polypeptides include Bcl-G_L and Bcl-G_S.

30 The invention also provides vectors containing Bcl-G nucleic acids, host cells containing such vectors,

Bcl-G anti-sense nucleic acids and related compositions. The invention additionally provides Bcl-G oligonucleotides that can be used to hybridize to or amplify a Bcl-G nucleic acid. Anti-Bcl-G specific antibodies are also provided. Further provided are kits containing Bcl-G nucleic acids or Bcl-G specific antibodies. Such kits and reagents can be used to diagnose cancer, monitor response to therapy, or predict the prognosis of a cancer patient. The invention additionally provides methods of modulating apoptosis using Bcl-G polypeptides, encoding nucleic acids, or compounds that modulate the activity or expression of Bcl-G polypeptides. The methods for modulating apoptosis can be used to treat diseases such as cancer.

15

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of human Bcl-G_L cDNA (SEQ ID NO:1).

Figure 2 shows the nucleotide sequence of the coding region of human Bcl-G_L cDNA (nucleotide 196-1179 of SEQ ID NO:1) and the encoded amino acid sequence (SEQ ID NO:2). Bcl-G_L contains a BH3 domain (²¹⁶LKYSGDQLE²²⁴; SEQ ID NO:5) and a BH2 domain (³⁰⁷PWQQHGGWE³¹⁶; SEQ ID NO:6).

20

Figure 3 shows the nucleotide sequence of human Bcl-G_S cDNA (SEQ ID NO:3).

25

Figure 4 shows the nucleotide sequence of the coding region of human Bcl-G_S cDNA (nucleotide 196-954 of SEQ ID NO:3) and the encoded amino acid sequence (SEQ ID NO:4). Bcl-G_S contains only the BH3 domain (²¹⁶LKYSGDQLE²²⁴; SEQ ID NO:5).

Figure 5 shows sequence analysis of Bcl-G cDNAs. Figure 5A shows the predicted amino acid sequences of the Bcl-G_L and Bcl-G_S proteins, with the BH2 and BH3 domains in bold-type and residue numbers indicated. The predicted proteins are identical from residues 1-226. The unique C-terminal region of Bcl-G_S is indicated in italics type. Figure 5B shows an alignment of the BH2 domains of Bcl-G_L (SEQ ID NO:9) and several other Bcl-2 family proteins (SEQ ID NOS:10-17, respectively). Identical and similar residues are shown in black and gray blocks, respectively. Figure 5C shows an alignment of the BH3 domains Bcl-G (SEQ ID NO:18) and several other Bcl-2 family proteins (SEQ ID NOS:19-26, respectively). Figure 5D shows the exon-intron organization of the *BCL-G* gene. The *BCL-G* gene contains 6 exons, spanning a ~30 kb region of chromosome 12. Alternative splicing at the 5'-end of exon 5 accounts for the production of the Bcl-G_L and Bcl-G_S proteins, where splice-acceptor sites at nucleotide positions 63,870 versus 63,797 in BAC clone RPCI 11-267J23 (GenBank AC007537) are utilized for Bcl-G_L and Bcl-G_S, respectively. The positions of the start and termination codons are indicated, with coding regions in gray blocks and non-coding 5'-UTR and 3'-UTR sequence in open blocks. The BH3 domain is located in exon 4 of both Bcl-G_L and Bcl-G_S, while the BH2 domain resides in exon 5 of Bcl-G_L.

Figure 6 shows mapping of Bcl-G to chromosome 12p12.3.

Figure 7 shows expression of Bcl-G_S and Bcl-G_L in human tissues. The expression of transcripts encoding Bcl-G_L or Bcl-G_S was examined by RT-PCR. First-strand cDNA prepared using RNA samples from various adult human tissues was PCR amplified using primers specific for Bcl-

G_L and Bcl- G_S , based on differences in splice-acceptor utilization in exon 5. The primers flank an intron in both cases, thus excluding amplification due to contaminating genomic DNA. PCR products were size-fractionated in 2% agarose gels, stained with ethidium bromide, then photographed under UV-illumination.

Figure 8 shows the effect of Bcl- G_S on cell death. PC-3 cells were transfected with pcDNA3.1/Myc/His (control), pcDNA3.1/Myc/His/Bcl- G_S (Bcl- G_S), pcDNA3.1/Myc/His/Bcl- G_S + pRC/CMV/Bcl-2 (Bcl- G_S + Bcl-2), pRC/CMV/Bax (Bax), or pRC/CMV/Bax + pRC/CMV/Bcl-2 (Bax + Bcl-2). Cells were tested for cell death 24 hours after transfection.

Figure 9 shows induction of apoptosis by Bcl-G. Figure 9A shows the results of transfecting plasmids encoding GFP, GFP-Bcl- G_S , or GFP-Bcl- G_L into Cos-7 cells alone or in combination with a plasmid encoding Bcl- X_L . Apoptosis was examined by DAPI staining at 24 h post-transfection (mean + SD; n = 3) (top). Levels of GFP and GFP-Bcl-G fusion proteins were examined by immunoblotting lysates from transfected Cos-7 cells (20 μ g per lane) and anti-GFP antibody with ECL-based detection (middle). Equal loading was confirmed by reprobing the same membrane with anti-Tubulin antibody (bottom). Figure 9B shows the results of transfecting plasmids encoding GFP, GFP-Bcl- G_S or the mutant proteins, Bcl- G_S (Δ BH3) and GFP-Bcl- G_S (L216E) into Cos-7 cells. The percentage of apoptotic cells was examined 1 day later as above (top). Protein expression was assessed by immunoblotting as above, using anti-GFP (middle) or anti-Tubulin (bottom) antibodies.

Figure 10 shows interactions of Bcl-G_s and Bcl-G_L with Bcl-X_L. 293T cells were transiently transfected with plasmids encoding GFP, GFP-Bcl-G_L, GFP-Bcl-G_s, GFP-Bcl-G_s (DBH3), or GFP-Bcl-G_s (L218E). Cells were lysed 1 day later and immunoprecipitations were performed using anti-GFP antibody. Immune-complexes (prepared from 2 mg lysate) (top) and lysates (20 µg protein) (bottom) were subjected to SDS-PAGE/immunoblot analysis using anti-Bcl-X_L (top) and anti-GFP (bottom) antibodies, respectively.

Figure 11 shows microscopic evaluation of intracellular distributions of Bcl-G_L and Bcl-G_s. Plasmids encoding GFP (A), GFP-Bcl-G_L (B), GFP-Bcl-G_s (C), and GFP-Bcl-G_s (ΔBH3) (D) were transfected into Cos-7 cells. Cells were fixed 1 day later and examined by confocal microscopy.

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there are provided nucleic acids encoding Bcl-G polypeptides, or functional polypeptide fragments thereof. As used herein, the term "Bcl-G" refers to sub-family members of the Bcl-2 family of proteins, wherein said Bcl-G comprises a BH3 domain (SEQ ID NOS:5 or 9). The human Bcl-G gene has been found to map to chromosome 12p12.3 (Example II). This region of chromosome 12 is frequently deleted in cancer cells, in particular in acute lymphoblastic leukemia (ALL) and other solid tumor cells (Baens et al., (1999) Genomics 56:40-50 (1999); Hatta et al., Br. J. Cancer 75:1256-1262 (1997); Kibel et al., Cancer Res. 58:5652-5655 (1998); Baccichet et al., Br. J. Haematol. 99:107-114 (1997); Aissani et al., Leuk. Lymphoma 34:231-239). This region is deleted in a subset of prostate (approximately 50%), ovarian (approximately

30%) and leukemias (approximately 30%). Therefore, Bcl-G can function as a tumor suppressor. Furthermore, the presence or absence of Bcl-G nucleic acid or polypeptide or changes in Bcl-G nucleic acid or polypeptide expression, can serve as a marker for predisposition or progression of cancer, for example, prostate, ovarian and leukemia. Thus, the invention Bcl-G nucleic acids and/or polypeptides can be used for screening for cancer and/or for developing drug candidates for the treatment of cancer. Invention Bcl-G nucleic acids and/or polypeptides can also be used for discovery of drugs, as disclosed herein, that suppress autoimmunity, inflammation, allergy, allograft rejection, sepsis, and other diseases, including inflammatory diseases.

A new member of the Bcl-2 family was identified, Bcl-G (see Examples). The human *BCL-G* gene consists of 6 exons, resides on chromosome 12p12, and encodes two proteins through alternative mRNA splicing: Bcl-G (long) and Bcl-G (short) consisting of 327 and 252 (length) amino acids, respectively. Bcl-G_L and Bcl-G_S are identical in their first 226 amino-acids but diverge thereafter. Among the Bcl-2 Homology (BH) domains previously recognized in Bcl-2 family proteins, the BH3 domain is found in both Bcl-G_L and Bcl-G_S, but only the longer Bcl-G_L protein possesses a BH2 domain. Bcl-G_L mRNA is expressed widely in normal human tissues, whereas Bcl-G_S mRNA was found only in testis. Over-expression of Bcl-G_L or Bcl-G_S in cells induced apoptosis, but Bcl-G_S was far more potent than Bcl-G_L. Apoptosis induction by Bcl-G_S depended on the BH3 domain, and was suppressed by co-expression of anti-apoptotic Bcl-XL protein. Bcl-XL also co-immunoprecipitated with Bcl-G_S but not with mutants of Bcl-G_S in which the BH3 domain was deleted or mutated and not with Bcl-G_L. Bcl-G_S was predominantly localized to

cytosolic organelles whereas Bcl-G_L was diffusely distributed throughout the cytosol. The findings suggest that Bcl-G_L is likely in a latent state, whereas the shorter Bcl-G_S protein is constitutively active.

5 The term "biologically active" or "functional", when used herein as a modifier of an invention Bcl-G, or polypeptide fragment thereof, refers to a polypeptide that exhibits functional characteristics similar to Bcl-G, including those disclosed herein (see Examples I-IX).
10 As disclosed herein, Bcl-G induces apoptosis (see Example IV). Therefore, one function of Bcl-G is a pro-apoptotic function. The pro-apoptotic function of Bcl-G is inhibited by co-expression of the anti-apoptotic protein Bcl-2 (Example IV). Therefore, another function of Bcl-G
15 is modulation by or interaction with an anti-apoptotic protein such as for example, Bcl-2 family member, including Bcl-2 or Bcl-X_L, and the like. Bcl-G can function to heterodimerize with a Bcl-2 family member, thereby modulating the apoptotic activity of Bcl-G and/or
20 the Bcl-2 family member. For example, the interaction of Bcl-G_S with Bcl-X_L was found to be BH3 domain dependent, and, thus, the pro-apoptotic activity of Bcl-G_S correlates with its ability to bind Bcl-X_L (see Example VI).

 Bcl-G is also contemplated herein as having the
25 ability to function as an ion channel. Additionally, Bcl-G is contemplated herein as having the ability to function target to mitochondria, for example, for example, by binding directly to mitochondria or via binding to a protein that is associated with mitochondria
30 such as Bcl-2 or Bcl-X_L. Bcl-G can also function to bind adenine nucleotide transporter (ANT) and to other proteins such as voltage-dependent anion channel (VDAC).

Because Bcl-G is located on chromosome 12 in a region that is frequently deleted in cancer cells (Example II) it is contemplated herein that Bcl-G functions as a tumor suppressor. Another functional activity of Bcl-G is the ability to act as an immunogen for the production of polyclonal and monoclonal antibodies that bind specifically to an invention Bcl-G. Thus, an invention nucleic acid encoding Bcl-G will encode a polypeptide specifically recognized by an antibody that also specifically recognizes the Bcl-G protein including the amino acid sequence, set forth in SEQ ID NOS:2, 4 or 42. Such immunologic activity can be assayed by any method known to those of skill in the art. Therefore, Bcl-G functional fragments include polypeptide fragments that function as immunogens for generating a Bcl-G-specific antibody and fragments that specifically bind to a Bcl-G-specific antibody.

Bcl-2 family proteins are central regulators of apoptosis (reviewed in Reed, J. C., Nature, 387:773-776 (1997); Adams & Cory, Science, 281:1322-1326 (1998); Gross et al., Genes Dev., 13:1899-1911 (1999)). Bcl-2 family proteins are conserved throughout the animal kingdom, with homologues identified in both vertebrates and invertebrates. These proteins contain up to four conserved Bcl-2 Homology (BH) domains, BH1, BH2, BH3, and BH4, which are recognized by their amino-acid sequence similarity. Both anti-apoptotic and pro-apoptotic Bcl-2 family proteins have been identified. These proteins control cell life-death decisions through their effects on events such as mitochondrial release of proteins involved in activation of caspase-family cell death proteases (reviewed in Gross et al., Genes Dev., 13:1899-1911 (1999); Green & Reed, Science, 281:1309-1312 (1998); Kroemer & Reed, Nature Medicine, 6:513-519 (2000)). Many

Bcl-2 family proteins are capable of physically interacting with each other, forming a complex network of homo-and heterodimers, and these physical interactions sometimes play important roles in the opposing effects of pro- and anti-apoptotic members of the family.

The pro-apoptotic members of the Bcl-2 family can be broadly classified into two groups. One group, including Bax, Bak, and Bok in humans, shares structural similarity with the pore-forming domains of certain bacterial toxins and is capable of forming pores in synthetic membranes *in vitro* (Schendel et al., Cell Death Differ., 5:372-380 (1998); Antonsson et al., Science, 277:370-372 (1997); Schlesinger et al., Proc. Natl. Acad. Sci. USA, 94:11357-11362 (1997); Shimizu et al., J Biol Chem., 16:12321-12325 (2000)). These protein exhibit cytotoxic effects independently of their ability to bind other Bcl-2 family proteins, including Bcl-2 and other cytoprotective members of the family such as Bcl-X_L, Bcl-W, Bfl-1, and Mcl-1. The second group of pro-apoptotic Bcl-2 family proteins varies widely in their amino-acid sequences, often containing only a single region of similarity, the BH3 domain. These "BH3-only" proteins appear to possess no intrinsic or autonomous cytodestructive activity, and instead operate as trans-dominant inhibitors of the survival proteins. Their antagonism of proteins such as Bcl-2 and Bcl-X_L depends on binding via their BH3 domains to a hydrophobic pocket on target anti-apoptotic proteins (Kelekar & Thompson, Trends Cell Biol., 8:324-330 (1998)).

Gene knock-out studies in mice have demonstrated non-redundant roles for various Bcl-2 family genes in regulating cell life and death in specific tissues or under particular physiological or pathological

circumstances (Veis et al., Cell, 75:229-240 (1993); Motoyama et al., Science, 267:1506-1510 (1995); Knudson et al., Science, 270:96-99 (1995); Bouillet et al., Science, 286:1735-8 (1999); Yin et al., Nature, 400:886-891 (1999)). Thus, it is important to identify all members of the Bcl-2 family and to delineate the cellular contexts in which they contribute to apoptosis regulation. As disclosed herein, a new member of the Bcl-2 family, Bcl-G, has been cloned and characterized.

The nucleic acid molecules described herein are useful for producing invention proteins, when such nucleic acids are incorporated into a variety of protein expression systems known to those of skill in the art. In addition, such nucleic acid molecules or fragments thereof can be labeled with a readily detectable substituent and used as hybridization probes for assaying for the presence and/or amount of an invention Bcl-G gene or mRNA transcript in a given sample. The nucleic acid molecules described herein, and fragments thereof, are also useful as primers and/or templates in a PCR reaction for amplifying genes encoding invention proteins described herein.

The term "nucleic acid", also referred to as polynucleotides, encompasses ribonucleic acid (RNA) or deoxyribonucleic acid (DNA), probes, oligonucleotides, and primers and can be single stranded or double stranded. DNA can be either complementary DNA (cDNA) or genomic DNA, and can represent the sense strand, the anti-sense strand or both. Examples of nucleic acids are RNA, cDNA, or isolated genomic DNA encoding an Bcl-G polypeptide. Such nucleic acids include, but are not limited to, nucleic acids comprising substantially the same nucleotide sequence as set forth in SEQ ID NOS:1, 3

or 41. In general, a genomic sequence of the invention includes regulatory regions such as promoters, enhancers, and introns that are outside of the exons encoding a Bcl-G but does not include proximal genes that do not encode Bcl-G.

Use of the terms "isolated" and/or "purified" in the present specification and claims as a modifier of DNA, RNA, polypeptides or proteins means that the DNA, RNA, polypeptides or proteins so designated have been produced in such form by the hand of man, and thus are separated from their native *in vivo* cellular environment.

As employed herein, the term "substantially the same nucleotide sequence" refers to DNA having sufficient identity to the reference polynucleotide, such that it will hybridize to the reference nucleotide under moderately stringent hybridization conditions. In one embodiment, DNA having substantially the same nucleotide sequence as the reference nucleotide sequence encodes substantially the same amino acid sequence as that set forth in any of SEQ ID NOS:2, 4 or 42. In another embodiment, DNA having "substantially the same nucleotide sequence" as the reference nucleotide sequence has at least 60% identity with respect to the reference nucleotide sequence. DNA having substantially the same nucleotide sequence can have at least 70%, at least 90%, or at least 95% identity to the reference nucleotide sequence.

As used herein, a "modification" of a nucleic acid can also include one or several nucleotide additions, deletions, or substitutions with respect to a reference sequence. A modification of a nucleic acid can include substitutions that do not change the encoded

amino acid sequence due to the degeneracy of the genetic code. Such modifications can correspond to variations that are made deliberately, or which occur as mutations during nucleic acid replication.

5 Exemplary modifications of the recited Bcl-G sequences include sequences that correspond to homologs of other species, including mammalian species such as mouse, primates, including monkey and baboon, rat, rabbit, bovine, porcine, ovine, canine, feline, or other
10 animal species. The corresponding Bcl-G sequences of non-human species can be determined by methods known in the art, such as by PCR or by screening genomic, cDNA or expression libraries.

Another exemplary modification of the invention
15 Bcl-G can correspond to splice variant forms of the Bcl-G nucleotide sequence. Additionally, a modification of a nucleotide sequence can include one or more non-native nucleotides, having, for example, modifications to the base, the sugar, or the phosphate portion, or having a
20 modified phosphodiester linkage. Such modifications can be advantageous in increasing the stability of the nucleic acid molecule.

Furthermore, a modification of a nucleotide sequence can include, for example, a detectable moiety,
25 such as a radiolabel, a fluorochrome, a ferromagnetic substance, a luminescent tag or a detectable binding agent such as biotin. Such modifications can be advantageous in applications where detection of a Bcl-G nucleic acid molecule is desired.

30 The invention also encompasses nucleic acids which differ from the nucleic acids shown in SEQ ID NOS:

1, 3 or 41, but which have the same phenotype.
Phenotypically similar nucleic acids are also referred to
as "functionally equivalent nucleic acids". As used
herein, the phrase "functionally equivalent nucleic
5 acids" encompasses nucleic acids characterized by slight
and non-consequential sequence variations that will
function in substantially the same manner to produce the
same protein product(s) as the nucleic acids disclosed
herein. In particular, functionally equivalent nucleic
10 acids encode polypeptides that are the same as those
encoded by the nucleic acids disclosed herein or that
have conservative amino acid variations. For example,
conservative variations include substitution of a non-
polar residue with another non-polar residue, or
15 substitution of a charged residue with a similarly
charged residue. These variations include those
recognized by skilled artisans as those that do not
substantially alter the tertiary structure of the
protein.

20 Further provided are nucleic acids encoding
Bcl-G polypeptides that, by virtue of the degeneracy of
the genetic code, do not necessarily hybridize to the
invention nucleic acids under specified hybridization
conditions. As used herein, the term "degenerate" refers
25 to codons that differ in at least one nucleotide from a
reference nucleic acid, but encode the same amino acids
as the reference nucleic acid. Nucleic acids encoding
the invention Bcl-G polypeptides can be comprised of
nucleotides that encode substantially the same amino acid
30 sequence as set forth in SEQ ID NOS:2, 4 or 42.

The invention provides an isolated nucleic acid
encoding a Bcl-G polypeptide, or a functional fragment
thereof. The invention also provides an isolated nucleic

acid encoding a Bcl-G polypeptide, or a functional fragment thereof, comprising a nucleic acid selected from:

- 5 (a) nucleic acid encoding the amino acid sequence set forth in SEQ ID NOS:2, 4 or 42, or
- 10 (b) nucleic acid that hybridizes to the nucleic acid of (a) under moderately stringent conditions, wherein said nucleic acid contiguously encodes biologically active Bcl-G, or
- (c) nucleic acid degenerate with respect to either (a) or (b) above, wherein said nucleic acid encodes biologically active Bcl-G.

15 In one embodiment, preferred Bcl-G polypeptide include a long form termed Bcl-G_L and a short form termed Bcl-G_S. Bcl-G_L contains a BH3 and a BH2 domain, whereas Bcl-G_S contains only a BH3 domain. Bcl-G_S has been found to possess pro-apoptotic activity similar to Bax (see Example III).

20 Hybridization refers to the binding of complementary strands of nucleic acid, for example, sense:antisense strands or probe:target-nucleic acid to each other through hydrogen bonds, similar to the bonds that naturally occur in chromosomal DNA. Stringency

25 levels used to hybridize a given probe with target-DNA can be readily varied by those of skill in the art.

30 The phrase "stringent hybridization" is used herein to refer to conditions under which polynucleic acid hybrids are stable. As known to those of skill in the art, the stability of hybrids is reflected in the

melting temperature (T_m) of the hybrids. In general, the stability of a hybrid is a function of sodium ion concentration and temperature. Typically, the hybridization reaction is performed under conditions of lower stringency, followed by washes of varying, but higher, stringency. Reference to hybridization stringency relates to such washing conditions.

As used herein, the phrase "moderately stringent hybridization" refers to conditions that permit target-nucleic acid to bind a complementary nucleic acid. The hybridized nucleic acids will generally have at least about 60% identity, at least about 75% identity, more at least about 85% identity; or at least about 90% identity. Moderately stringent conditions are conditions equivalent to hybridization in 50% formamide, 5X Denhart's solution, 5X SSPE, 0.2% SDS at 42°C, followed by washing in 0.2X SSPE, 0.2% SDS, at 42°C.

The phrase "high stringency hybridization" refers to conditions that permit hybridization of only those nucleic acid sequences that form stable hybrids in 0.018M NaCl at 65°C, for example, if a hybrid is not stable in 0.018M NaCl at 65°C, it will not be stable under high stringency conditions, as contemplated herein. High stringency conditions can be provided, for example, by hybridization in 50% formamide, 5X Denhart's solution, 5X SSPE, 0.2% SDS at 42°C, followed by washing in 0.1X SSPE, and 0.1% SDS at 65°C.

The phrase "low stringency hybridization" refers to conditions equivalent to hybridization in 10% formamide, 5X Denhart's solution, 6X SSPE, 0.2% SDS at 22°C, followed by washing in 1X SSPE, 0.2% SDS, at 37°C. Denhart's solution contains 1% Ficoll, 1%

polyvinylpyrrolidone, and 1% bovine serum albumin (BSA).
20X SSPE (sodium chloride, sodium phosphate, ethylene
diamide tetraacetic acid (EDTA)) contains 3M sodium
chloride, 0.2M sodium phosphate, and 0.025 M (EDTA).
5 Other suitable moderate stringency and high stringency
hybridization buffers and conditions are well known to
those of skill in the art and are described, for example,
in Sambrook et al., Molecular Cloning: A Laboratory
Manual, 2nd ed., Cold Spring Harbor Press, Plainview, New
10 York (1989); and Ausubel et al., *supra*, 1999). Nucleic
acids encoding polypeptides hybridize under moderately
stringent or high stringency conditions to substantially
the entire sequence, or substantial portions, for
example, typically at least 15-30 nucleotides of the
15 nucleic acid sequence set forth in SEQ ID NOS:1, 3 or 41.

The invention also provides a modification of a
Bcl-G nucleotide sequence that hybridizes to a Bcl-G
nucleic acid molecule, for example, a nucleic acid
molecule referenced as SEQ ID NOS:1, 3 or 41, under
20 moderately stringent conditions. Modifications of Bcl-G
nucleotide sequences, where the modification has at least
60% identity to a Bcl-G nucleotide sequence, are also
provided. The invention also provides modification of a
Bcl-G nucleotide sequence having at least 65% identity,
25 at least 70% identity, at least 75% identity, at least
80% identity, at least 85% identity, at least 90%
identity, or at least 95% identity.

Identity of any two nucleic acid sequences can
be determined by those skilled in the art based, for
30 example, on a BLAST 2.0 computer alignment, using default
parameters. BLAST 2.0 searching is available at
<http://www.ncbi.nlm.nih.gov/gorf/bl2.html>., as described
by Tatiana et al., FEMS Microbiol Lett. 174:247-250

(1999); Altschul et al., Nucleic Acids Res., 25:3389-3402 (1997).

One means of isolating a nucleic acid encoding a Bcl-G polypeptide is to probe a cDNA library or genomic library with a natural or artificially designed nucleic acid probe using methods well known in the art. Nucleic acid probes derived from the Bcl-G gene are particularly useful for this purpose. DNA and cDNA molecules that encode Bcl-G polypeptides can be used to obtain complementary genomic DNA, cDNA or RNA from mammals, for example, human, mouse, rat, rabbit, pig, and the like, or other animal sources, or to isolate related cDNA or genomic clones by the screening of cDNA or genomic libraries, by methods well known in the art (see, for example, Sambrook et al., supra, 1989; Ausubel et al., supra, 1999).

The invention additionally provides a nucleic acid that hybridizes under high stringency conditions to the Bcl-G coding portion of any of SEQ ID NOS:1, 3 or 41. The invention also provides a nucleic acid having a nucleotide sequence the same or substantially the same as set forth in any of SEQ ID NOS:1, 3 or 41.

The invention also provides a method for identifying nucleic acids encoding a mammalian Bcl-G by contacting a sample containing nucleic acids with one or more Bcl-G oligonucleotides, wherein the contacting is effected under high stringency hybridization conditions, and identifying a nucleic acid that hybridizes to the oligonucleotide. The invention additionally provides a method of detecting a Bcl-G nucleic acid molecule in a sample by contacting the sample with two or more Bcl-G

oligonucleotides, amplifying a nucleic acid molecule, and detecting the amplification. The amplification can be performed, for example, using PCR. The invention further provides oligonucleotides that function as single
5 stranded nucleic acid primers for amplification of a Bcl-G nucleic acid, wherein the primers comprise a nucleic acid sequence derived from the nucleic acid sequences set forth as SEQ ID NOS:1, 3 or 41.

In accordance with a further embodiment of the
10 present invention, optionally labeled Bcl-G-encoding nucleic acids, or fragments thereof, can be employed to probe a library, for example, a cDNA or genomic library, and the like for additional nucleic acid sequences encoding novel Bcl-G polypeptides. Construction of
15 suitable cDNA libraries is well-known in the art. Screening of such a cDNA library is initially carried out under low-stringency conditions, which comprise a temperature of less than about 42°C, a formamide concentration of less than about 50%, and a moderate to
20 low salt concentration.

Presently preferred probe-based screening conditions comprise a temperature of about 37°C, a formamide concentration of about 20%, and a salt concentration of about 5X sodium chloride, sodium citrate
25 (SSC; 20X SSC contains 3M sodium chloride, 0.3M sodium citrate, pH 7.0). Such conditions will allow the identification of sequences having a substantial degree of similarity with the probe sequence, without requiring perfect identity. The phrase "substantial similarity"
30 refers to sequences which share at least 50% identity. Hybridization conditions are selected which allow the identification of sequences having at least 70% identity with the probe, while discriminating against sequences

having a lower degree of identity with the probe. As a result, nucleic acids having substantially the same nucleotide sequence as SEQ ID NOS:1, 3 or 41 are obtained.

5 As used herein, a nucleic acid "probe" is single-stranded nucleic acid, or analogs thereof, that has a sequence of nucleotides that includes at least 14, at least 20, at least 50, at least 100, at least 200, at least 300, at least 400, or at least 500 contiguous bases
10 that are the same as or the complement thereof, any contiguous bases set forth in any of SEQ ID NOS:1, 3 or 41. In addition, the entire cDNA encoding region of an invention Bcl-G, or the entire sequence corresponding to SEQ ID NOS:1, 3 or 41 can be used as a probe. Probes can
15 be labeled by methods well-known in the art, as described hereinafter, and used, for example, in various diagnostic kits.

 The invention additionally provides a Bcl-G oligonucleotide comprising between 15 and 300 contiguous
20 nucleotides of SEQ ID NOS:1, 3 or 41, or the anti-sense strand thereof. As used herein, the term "oligonucleotide" refers to a nucleic acid molecule that includes at least 15 contiguous nucleotides from a reference nucleotide sequence, can include at least 16,
25 17, 18, 19, 20 or at least 25 contiguous nucleotides, and often includes at least 30, 40, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, up to 350 contiguous nucleotides from the reference nucleotide sequence. The reference nucleotide sequence can be the
30 sense strand or the anti-sense strand.

 The Bcl-G oligonucleotides of the invention that contain at least 15 contiguous nucleotides of a

reference Bcl-G nucleotide sequence are able to hybridize to Bcl-G under moderately stringent hybridization conditions and thus can be advantageously used, for example, as probes to detect Bcl-G DNA or RNA in a sample, and to detect splice variants thereof; as sequencing or PCR primers; as antisense reagents to block transcription of Bcl-G RNA in cells; or in other applications known to those skilled in the art in which hybridization to a Bcl-G nucleic acid molecule is desirable.

It is understood that a Bcl-G nucleic acid molecule, as used herein, specifically excludes previously known nucleic acid molecules consisting of nucleotide sequences having identity with the Bcl-G nucleotide sequence (SEQ ID NOS:1, 3 or 41), such as Expressed Sequence Tags (ESTs), Sequence Tagged Sites (STSs) and genomic fragments, deposited in public databases such as the nr, dbest, dbsts, gss and htgs databases, which are available for searching at <http://www.ncbi.nlm.nih.gov/blast/blast.cgi?Jform=0>, using the program BLASTN 2.0.9 described by Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997).

In particular, a Bcl-G nucleic acid molecule specifically excludes nucleic acid molecules consisting of any of the nucleotide sequences having the Genbank (gb), EMBL (emb) or DDBJ (dbj) accession numbers described below. Similarly, a Bcl-G polypeptide fragment specifically excludes the amino acid fragments encoded by the nucleotide sequences having the GenBank accession numbers described below. GenBank accession numbers specifically excluded include AC005903, AC007439, AW000827, AA399486, AW001213, AI478889, AA400686, AA398276, AI240211, and AA536718. The human BAC

referenced as GenBank accession No. AC007537 is also specifically excluded from a Bcl-G nucleic acid.

5 The isolated Bcl-G nucleic acid molecules of the invention can be used in a variety of diagnostic and therapeutic applications. For example, the isolated Bcl-G nucleic acid molecules of the invention can be used as probes, as described above; as templates for the recombinant expression of Bcl-G polypeptides; or in screening assays such as two-hybrid assays to identify
10 cellular molecules that bind Bcl-G.

 Another useful method for producing a Bcl-G nucleic acid molecule of the invention involves amplification of the nucleic acid molecule using PCR and Bcl-G oligonucleotides and, optionally, purification of
15 the resulting product by gel electrophoresis. Either PCR or RT-PCR can be used to produce a Bcl-G nucleic acid molecule having any desired nucleotide boundaries. Desired modifications to the nucleic acid sequence can also be introduced by choosing an appropriate
20 oligonucleotide primer with one or more additions, deletions or substitutions. Such nucleic acid molecules can be amplified exponentially starting from as little as a single gene or mRNA copy, from any cell, tissue or species of interest.

25 The invention thus provides methods for detecting Bcl-G nucleic acid in a sample. The methods of detecting Bcl-G nucleic acid in a sample can be either qualitative or quantitative, as desired. For example, the presence, abundance, integrity or structure of a Bcl-G
30 can be determined, as desired, depending on the assay format and the probe used for hybridization or primer pair chosen for application.

Useful assays for detecting Bcl-G nucleic acid based on specific hybridization with an isolated Bcl-G nucleic acid molecule are well known in the art and include, for example, *in situ* hybridization, which can be used to detect altered chromosomal location of the nucleic acid molecule, altered gene copy number, and RNA abundance, depending on the assay format used. Other hybridization assays include, for example, Northern blots and RNase protection assays, which can be used to determine the abundance and integrity of different RNA splice variants, and Southern blots, which can be used to determine the copy number and integrity of DNA. A Bcl-G hybridization probe can be labeled with any suitable detectable moiety, such as a radioisotope, fluorochrome, chemiluminescent marker, biotin, or other detectable moiety known in the art that is detectable by analytical methods.

Useful assays for detecting a Bcl-G nucleic acid in a sample based on amplifying a Bcl-G nucleic acid with two or more Bcl-G oligonucleotides are also well known in the art, and include, for example, qualitative or quantitative polymerase chain reaction (PCR); reverse-transcription PCR (RT-PCR); single strand conformational polymorphism (SSCP) analysis, which can readily identify a single point mutation in DNA based on differences in the secondary structure of single-strand DNA that produce an altered electrophoretic mobility upon non-denaturing gel electrophoresis; and coupled PCR, transcription and translation assays, such as a protein truncation test, in which a mutation in DNA is determined by an altered protein product on an electrophoresis gel. Additionally, the amplified Bcl-G nucleic acid can be sequenced to detect mutations and mutational hot-spots, and specific assays for large-scale screening of samples to identify

such mutations can be developed.

The invention further provides an isolated Bcl-G polypeptide, or a functional fragment thereof, encoded by a Bcl-G nucleic acid of the invention. For example, the invention provides a polypeptide comprising the same or substantially the same amino acid sequence as Bcl-G_L (SEQ ID NO:2) or Bcl-G_S (SEQ ID NO:4). Also provided is a Bcl-G polypeptide encoded by a nucleotide sequence comprising the same or substantially the same nucleotide sequence as set forth in SEQ ID NOs:1 or 3. Additionally provided is mouse Bcl-G nucleotide sequence set forth as SEQ ID NO:41 (see Example IX).

Described herein is a new member of the BCL-2 gene family in humans, BCL-G (see Examples I-VIII). The BCL-G gene potentially encodes two protein products, Bcl-G_L and Bcl-G_S. Bcl-2 family proteins contain up to four conserved BH domains. The shorter Bcl-G_S protein contains only the BH3 domains, similar to several other pro-apoptotic Bcl-2 family proteins, including Bad, Hrk, Bik, Bim, Apr, and Egl1 (reviewed in Kelekar & Thompson, Trends Cell Biol., 8:324-330 (1998); Reed, J. Oncogene, 17:3225-3236 (1998)). In contrast, the longer Bcl-G_L protein contains a BH2 and BH3 domain. No other examples of Bcl-2 family proteins are known which combine BH2 and BH3 domain in the absence of BH1. Though the Bad protein was originally suggested to contain a BH2 domain (Yang et al., Blood, 84(Suppl.1):373a-380a (1994)), and has been shown to possess the BH3 domain, inspection of the BH2 region reveals very little similarity of amino-acid sequence with (Ottillie et al., J. Biol. Chem., 272:30866-30872 (1997)) other BH2 domains. In contrast, the BH2 of Bcl-G_L contains a stretch of 8 of 8 residues showing identity or conservative amino-acid substitutions with

the BH2 domains of other family members. By comparison, the Bad sequence reveals only 3 of 8 identical or similar amino-acids in the same region. Thus, Bcl-G_L defines a novel structural variant within the Bcl-2 family of apoptosis-regulating proteins.

The production of different protein isoforms by alternative mRNA splicing is a common feature of *BCL-2* family genes, including *BCL-2*, *Bcl-X*, *MCL-1*, *BAX*, and *BIM* (Tsujiimoto & Croce, Proc. Natl. Acad. Sci. USA, 83:5214-5218 (1986); O'Connor et al., EMBO J., 17:384-395 (1998); Boise et al., Cell, 74:597-608 (1993); Oltvai et al., Cell, 74:609-619 (1993); Bingle et al., J. Biol. Chem., 275:22136-22146 (2000)). Unlike *BCL-X*, which encodes a longer and short protein, Bcl-X_L and Bcl-X_S, possessing anti-apoptotic and pro-apoptotic functions, respectively, the longer isoform of Bcl-G did not display anti-apoptotic activity. When over-expressed, Bcl-G_L induced modest and variable increases in apoptosis, whereas the shorter Bcl-G_S protein consistently exhibited potent cytotoxic activity. This behavior is reminiscent of the proteins encoded by the *BIM* gene, which include Bim-short (Bim_S), Bim-long (Bim_L) and Bim-Extra-Long (Bim_{EL}) (O'Connor et al., EMBO J., 17:384-395 (1998)). The longer proteins, Bim_L and Bim_{EL}, are sequestered in complexes with dynein light-chain (DLC) in association with microtubules, thus preventing them from interacting with target proteins such as Bcl-X_L on the surface of mitochondria and other organelles (Puthalakath et al., Mol. Cell, 3:287-96 (1999)). In contrast, because the shortest isoform, Bim_S, does not associate with DLC, it is free to interact with Bcl-X_L, Bcl-2, and other survival proteins and hence displays far more potent apoptotic activity when over-expressed in cells. By analogy, the longer Bcl-G_L protein could be sequestered in an inactive

complex with an unidentified protein.

Besides interactions with sequestering proteins, the activity of pro-apoptotic Bcl-2 family proteins can be suppressed by other mechanisms, including post-translational modifications. For example, the Bad protein is inactivated by phosphorylation. This protein can be directly or indirectly phosphorylated by several protein kinases, including PKA, PKB (Akt), Raf1, and Pak1, thus preventing it from dimerizing with target proteins such as Bcl-2 and Bcl-X_L (reviewed in Reed, J. Oncogene, 17:3225-3236 (1998); Datta et al., Genes Dev., 13:2905-2927 (1999)). The intracellular location of Bad varies, depending on its phosphorylation state, with phosphorylated Bad residing in the cytosol and unphosphorylated Bad associated with mitochondria and other intracellular organelles where Bcl-2 and Bcl-X_L are located. In this regard, the Bcl-G_L protein contains candidate phosphorylation sites for protein kinase A (PKA) and protein kinase C (PKC), including some not found in Bcl-G_S. However, *in vivo* phosphorylation of Bcl-G_L has not been observed in pilot experiments.

Another post-translational modification shown previously to activate latent pro-apoptotic Bcl-2 family proteins is proteolysis. Specifically, the Bid protein contains a N-terminal domain of ~56 amino-acids that masks its BH3 domain, reducing its ability to dimerize with other Bcl-2 family proteins. Upon cleavage by caspases, however, removal of the N-terminal domain exposes the BH3 domain and is associated with translocation of Bid from the cytosol to mitochondria, where it induces cytochrome c release and apoptosis (Li et al., Cell, 94:491-501 (1998); Luo et al., Cell, 94:481-490 (1998)). While Bcl-G_L contains candidate

caspase recognition sites, no significant cleavage of Bid has been observed *in vitro* using purified active caspases or in cells during apoptosis. It is possible, however, that a specific caspase not yet tested is capable of
5 cleaving and activating Bcl-G_L.

Though possessing no hydrophobic region that might anchor it in membranes, the Bcl-G_s protein was constitutively associated with intracellular organelles. Interestingly, removal of the BH3 domain did not
10 interfere with organellar-targeting of Bcl-G_s, but did abolish dimerization with Bcl-X_L. Thus, the BH3 domain apparently is not responsible for association of Bcl-G_s with intracellular organelles. This BH3-independent targeting of Bcl-G_s differs from some other "BH3-only"
15 Bcl-2 family proteins such as Bad, where it has been observed that removal of the BH3 domain abrogates binding to anti-apoptotic Bcl-2 family proteins as well as association with mitochondria (Zha et al., J. Biol. Chem., 272:24101-24104 (1997)).

20 The *BCL-G* gene resides on chromosome 12p12, a region deleted in ~50% of prostate cancers, ~30% of ovarian cancers, and ~30% of childhood acute lymphocytic leukemias (ALLs) (Kibel et al., J Urol., 1:192-196 (2000); Aissani et al., Leuk Lymphoma, 34:231-239 (1999);
25 Hatta et al., Br J Cancer, 75:1256-1262 (1997)). Given that at least one of the protein products of the *BCL-G* gene exhibits pro-apoptotic function, it is possible that *BCL-G* represents a tumor suppressor gene. However, thus far, somatic mutations in the exons of *BCL-G* have not
30 been detected nor evidence of deletion of both *BCL-G* alleles in tumor cell lines or primary tumor specimens tested so far. Further studies are required therefore to

determine whether loss of *BCL-G* expression occurs in tumors by means other than somatic alterations in gene structure and DNA sequence, such as changes in gene methylation or aberrant transcriptional or post-transcriptional regulation.

Investigation of the tissue-distribution of *Bcl-G_L* and *Bcl-G_S* mRNAs by RT-PCR revealed that *Bcl-G_L* mRNA is found in several normal adult tissues, whereas *Bcl-G_S* was detected only in testis. This finding indicates tissue-specific regulation of *Bcl-G_S* mRNA splicing. Tissue-specific splicing of other *Bcl-2* family mRNAs has been observed previously. For example, *Bcl-X* mRNA splicing events which generate the pro-apoptotic *Bcl-X_S* protein occur in the thymus during T-cell ontogeny and in the mammary gland during post-lactation involution, in association with extensive apoptosis induction (Boise et al., Cell, 74:597-608 (1993); Heermeier et al., Mech. Dev., 56:197-207 (1996)). Additional studies are performed to assess differential mRNA splicing patterns of *Bcl-G* transcripts during fetal development and following various scenarios in the adult where apoptosis occurs as part of a normal physiological response or an abnormal pathological reaction to environmental insults.

As employed herein, the term "substantially the same amino acid sequence" refers to amino acid sequences having at least about 70% identity with respect to the reference amino acid sequence, and retaining comparable functional and biological activity characteristic of the protein defined by the reference amino acid sequence. Preferably, proteins having "substantially the same amino acid sequence" will have at least about 80%, more preferably 90% amino acid identity with respect to the

reference amino acid sequence; with greater than about 95% amino acid sequence identity being especially preferred. It is recognized, however, that polypeptides, or encoding nucleic acids, containing less than the described levels of sequence identity arising as splice variants or that are modified by conservative amino acid substitutions, or by substitution of degenerate codons are also encompassed within the scope of the present invention.

Also encompassed by the term Bcl-G are functional fragments or polypeptide analogs thereof. The term "functional fragment" refers to a peptide fragment that is a portion of a full length Bcl-G protein, provided that the portion has a biological activity, as defined herein, that is characteristic of the corresponding full length protein. Thus, the invention also provides functional fragments of invention Bcl-G proteins, which can be identified using the binding and routine methods, such as bioassays described herein. A Bcl-G polypeptide functional fragment can be a BH3 or BH2 domain, for example, a BH3 domain referenced as SEQ ID NOS:5 or 9 or a BH2 domain referenced as SEQ ID NOS:6 or 18. The BH3 domain of Bcl-G is 33% identical to the BH3 domain of Bcl-2, 44% identical to the BH3 domain of Bcl-X_L, and 66% identical to the BH3 domain of Bax.

In addition, a functional fragment of a Bcl-G polypeptide can be Bax homology region. A region upstream of the BH3 domain shares a high degree of homology with Bax, including a 12 amino acid residue motif that is 70% identical between Bcl-G and Bax. Therefore, such a Bax homology region can function similarly to Bax, for example, as a possible binding domain. The N-terminal 150 amino acids of Bcl-G are not

similar to any known amino acid sequence available in public databases. Therefore, the N-terminal region of Bcl-G can function as Bcl-G-specific functional domain that confers a biological activity that is specific for Bcl-G relative to other members of the Bcl-2 family.

The invention also provides a chimeric protein comprising a domain selected from the group consisting of BH3 (SEQ ID NOS:5 or 9) and BH2 (SEQ ID NOS:6 or 18). A chimeric protein comprising a Bcl-G functional domain can be generated, for example, by recombinantly expressing a Bcl-G domain such as BH2 or BH3 fused to another polypeptide. Alternatively, the Bcl-G functional domain can be expressed as a fusion to another polypeptide.

In another embodiment of the invention, Bcl-G-containing chimeric proteins are provided comprising an invention Bcl-G, or fragments thereof, having the sequence of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:42, and further comprising one or more sequences from a heterologous protein. Sequences from heterologous proteins with which the Bcl-G or functional fragment thereof are fused can include, for example, glutathione-S-transferase, an antibody, or other proteins or functional fragments thereof which facilitate recovery of the chimera. Further proteins with which the Bcl-G or functional fragment thereof are fused will include, for example, luciferase, green fluorescent protein, an antibody, or other proteins or functional fragments thereof which facilitate identification of the chimera. Still further proteins with which the Bcl-G or functional fragment thereof are fused will include, for example, the LexA DNA binding domain, ricin, α -sarcin, an antibody, or other proteins which have therapeutic properties or other biological activity.

As such chimeric proteins include sequences from two different proteins, the resultant amino acid sequence of the chimeric protein will typically be a non-naturally occurring sequence. Thus, in accordance with this embodiment of the invention, there are provided chimeric proteins comprising an invention Bcl-G, or fragments thereof, having the sequence of SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:42, or a fragment thereof, provided the sequence of the chimeric protein is not naturally occurring.

In another embodiment of the invention, there are provided hetero-oligomers comprising invention Bcl-G polypeptides and fragments thereof, invention Bcl-G-containing proteins, Bcl-G-containing chimeric proteins, or combinations thereof. As disclosed herein, Bcl-G contains a BH3 domain, which functions as a ligand to bind Bcl-2 family members (Example I). Bcl-G can function to bind Bcl-2 family members. Thus, hetero-oligomers comprising invention Bcl-G polypeptides (SEQ ID NOS:2, 4 or 42) and fragments thereof, invention Bcl-G-containing proteins, Bcl-G-containing chimeric proteins, or combinations thereof, and further comprising Bcl-2 family members such as Bcl-2, Bcl-X_L or other Bcl-2 family members are provided.

As used herein, the term "polypeptide" when used in reference to Bcl-G is intended to refer to a peptide or polypeptide of two or more amino acids. The term "polypeptide analog" includes any polypeptide having an amino acid residue sequence substantially the same as a sequence specifically described herein in which one or more residues have been conservatively substituted with a functionally similar residue and which displays the ability to functionally mimic a Bcl-G as described

herein. A "modification" of a Bcl-G polypeptide also encompasses conservative substitutions of a Bcl-G polypeptide amino acid sequence. Conservative substitutions of encoded amino acids include, for
5 example, amino acids that belong within the following groups: (1) non-polar amino acids (Gly, Ala, Val, Leu, and Ile); (2) polar neutral amino acids (Cys, Met, Ser, Thr, Asn, and Gln); (3) polar acidic amino acids (Asp and Glu); (4) polar basic amino acids (Lys, Arg and His); and
10 (5) aromatic amino acids (Phe, Trp, Tyr, and His). Other minor modifications are included within Bcl-G polypeptides so long as the polypeptide retains some or all of its function as described herein.

The amino acid length of functional fragments
15 or polypeptide analogs of the present invention can range from about 5 amino acids up to the full-length protein sequence of an invention Bcl-G. In certain embodiments, the amino acid lengths include, for example, at least
20 about 10 amino acids, at least about 15, at least about 20, at least about 25, at least about 30, at least about 35, at least about 40, at least about 45, at least about 50, at least about 75, at least about 100, at least about 150, at least about 200, at least about 250 or more amino acids in length up to the full-length Bcl-G protein
25 sequence. The functional fragments can be contiguous amino acid sequences of a Bcl-G polypeptide, including contiguous amino acid sequences of SEQ ID NOS:2, 4 or 42.

A modification of a polypeptide can also include derivatives, analogues and functional mimetics
30 thereof, provided that such polypeptide displays the Bcl-G biological activity. For example, derivatives can include chemical modifications of the polypeptide such as alkylation, acylation, carbamylation, iodination, or any

modification that derivatizes the polypeptide. Such derivatized molecules include, for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups can be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups can be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine can be derivatized to form N-im-benzylhistidine. Also included as derivatives or analogues are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids, for example, 4-hydroxyproline, 5-hydroxylysine, 3-methylhistidine, homoserine, ornithine or carboxyglutamate, and can include amino acids that are not linked by peptide bonds. Polypeptides of the present invention also include any polypeptide having one or more additions and/or deletions of residues, relative to the sequence of a polypeptide whose sequence is shown herein, so long as Bcl-G activity is maintained.

A modification of a Bcl-G polypeptide includes functional mimetics thereof. Mimetics encompass chemicals containing chemical moieties that mimic the function of the polypeptide. For example, if a polypeptide contains two charged chemical moieties having functional activity, a mimetic places two charged chemical moieties in a spatial orientation and constrained structure so that the charged chemical function is maintained in three-dimensional space. Thus, a mimetic, which orients functional groups that provide a function of Bcl-G, are included within the meaning of a Bcl-G derivative. All of these modifications are

included within the term "polypeptide" so long as the Bcl-G polypeptide or functional fragment retains its function.

The invention provides an isolated Bcl-G polypeptide, or functional fragment thereof. The invention Bcl-G polypeptides can be isolated by a variety of methods well-known in the art, for example, recombinant expression systems described herein, precipitation, gel filtration, ion-exchange, reverse-phase and affinity chromatography, and the like. Other well-known methods are described in Deutscher et al., Guide to Protein Purification: Methods in Enzymology Vol. 182, (Academic Press, (1990)). Alternatively, the isolated polypeptides of the present invention can be obtained using well-known recombinant methods (see, for example, Sambrook et al., *supra*, 1989; Ausubel et al., *supra*, 1999). The methods and conditions for biochemical purification of a polypeptide of the invention can be chosen by those skilled in the art, and purification monitored, for example, by an immunological assay or a functional assay.

An example of the means for preparing the invention polypeptide(s) is to express nucleic acids encoding Bcl-G in a suitable host cell, such as a bacterial cell, a yeast cell, an amphibian cell such as an oocyte, or a mammalian cell, using methods well known in the art, and recovering the expressed polypeptide, again using well-known purification methods, so described herein. Invention polypeptides can be isolated directly from cells that have been transformed with expression vectors as described herein. Recombinantly expressed polypeptides of the invention can also be expressed as fusion proteins with appropriate affinity tags, such as

glutathione S transferase (GST) or poly His, and affinity purified. The invention polypeptide, biologically functional fragments, and functional equivalents thereof can also be produced by chemical synthesis. For example, 5 synthetic polypeptides can be produced using Applied Biosystems, Inc. Model 430A or 431A automatic peptide synthesizer (Foster City, CA) employing the chemistry provided by the manufacturer.

Bcl-G polypeptides can be administered to an 10 individual to increase an activity associated with a Bcl-G polypeptide, including induction of apoptosis or functioning as a tumor suppressor. For example, a Bcl-G polypeptide can be administered therapeutically to an individual using expression vectors containing nucleic 15 acids encoding Bcl-G polypeptides, as described below. In addition, Bcl-G polypeptides, or a functional portion thereof, can be directly administered to an individual. Methods of administering therapeutic polypeptides are well known to those skilled in the art, for example, as a 20 pharmaceutical composition.

In a particular embodiment, a Bcl-G polypeptide, or functional fragment thereof, can be administered to an individual so that the Bcl-G polypeptide or functional fragment is targeted to a tumor 25 to induce apoptosis or otherwise function as a tumor suppressor. One method of delivering a Bcl-G polypeptide to an intracellular target is to fuse a Bcl-G polypeptide or functional fragment to an intracellular-targeting peptide that can penetrate the cell membrane or otherwise 30 deliver a polypeptide to the intracellular environment such as via internalization, thereby causing the fused Bcl-G polypeptide to enter the cell. One example of such an intracellular-targeting peptides is a fusion to the

transduction domain of HIV TAT, which allows transduction of up to 100% of cells (Schwarze et al., Science 285:1569-1572 (1999); Vocero-Akbani et al., Nature Med. 5:29-33 (1999)).

5 Another example of such an intracellular-targeting peptide is the Antennapeida homeoprotein internalization domain (Holinger et al., J. Biol. Chem. 274:13298-13304 (1999)). Still another intracellular-targeting peptide is a peptide that is specific for a
10 cell surface receptor, which allows binding and internalization of a fusion polypeptide via receptor-mediated endocytosis (Ellerby et al., Nature Med. 5:1032-1038 (1999)). Such intracellular-targeting peptides that mediate specific receptor interactions can be
15 advantageously used to target a tumor (see Ellerby et al., *supra*, 1999). Alternatively, a Bcl-G polypeptide of the invention can be incorporated, if desired, into liposomes, microspheres or other polymer matrices (Gregoriadis, Liposome Technology, Vols. I to III,
20 2nd ed., CRC Press, Boca Raton FL (1993)).

 The invention additionally provides a method for modulating the activity of an oncogenic polypeptide by contacting the oncogenic polypeptide with a
25 substantially pure Bcl-G, or an oncogenic protein-binding fragment thereof. Bcl-G can function to bind oncogenic proteins such as Bcl-2. Therefore, Bcl-G or functional fragments that bind to an oncogenic protein such as Bcl-2 can be used to modulate the activity of the oncogenic protein.

30 The present invention also provides compositions containing an acceptable carrier and any of an isolated, purified Bcl-G mature protein or functional

polypeptide fragments thereof, alone or in combination with each other. These polypeptides or proteins can be recombinantly derived, chemically synthesized or purified from native sources. As used herein, the term

5 "acceptable carrier" encompasses any of the standard pharmaceutical carriers, such as phosphate buffered saline solution, water and emulsions such as an oil and water emulsion, and various types of wetting agents.

The invention thus provides a therapeutic

10 composition comprising a pharmaceutically acceptable carrier and a compound selected from the group consisting of a Bcl-G polypeptide, a functional fragment of Bcl-G, a Bcl-G modulating compound, and an anti-Bcl-G antibody. The invention additionally provides a method of treating

15 a pathology characterized by abnormal cell proliferation by administering an effective amount of the composition containing a pharmaceutically acceptable carrier and a compound selected from the group consisting of a Bcl-G polypeptide, a functional fragment of Bcl-G, a Bcl-G

20 modulating compound, and an anti-Bcl-G antibody.

Also provided are antisense-nucleic acids having a sequence capable of binding specifically with full-length or any portion of an mRNA that encodes Bcl-G polypeptides so as to prevent translation of the mRNA.

25 The antisense-nucleic acid can have a sequence capable of binding specifically with any portion of the sequence of the cDNA encoding Bcl-G polypeptides. As used herein, the phrase "binding specifically" encompasses the ability of a nucleic acid sequence to recognize a complementary

30 nucleic acid sequence and to form double-helical segments therewith via the formation of hydrogen bonds between the complementary base pairs. An example of an antisense-nucleic acid is an antisense-nucleic acid comprising

chemical analogs of nucleotides.

The present invention provides means to modulate levels of expression of Bcl-G polypeptides by recombinantly expressing Bcl-G anti-sense nucleic acids or employing synthetic anti-sense nucleic acid compositions (hereinafter SANC) that inhibit translation of mRNA encoding these polypeptides. Synthetic oligonucleotides, or other antisense-nucleic acid chemical structures designed to recognize and selectively bind to mRNA are constructed to be complementary to full-length or portions of an Bcl-G coding strand, including nucleotide sequences set forth in SEQ ID NOS:1, 3 or 41.

The SANC is designed to be stable in the blood stream for administration to a subject by injection, or in laboratory cell culture conditions. The SANC is designed to be capable of passing through the cell membrane in order to enter the cytoplasm of the cell by virtue of physical and chemical properties of the SANC, which render it capable of passing through cell membranes, for example, by designing small, hydrophobic SANC chemical structures, or by virtue of specific transport systems in the cell which recognize and transport the SANC into the cell. In addition, the SANC can be designed for administration only to certain selected cell populations by targeting the SANC to be recognized by specific cellular uptake mechanisms which bind and take up the SANC only within select cell populations. In a particular embodiment the SANC is an antisense oligonucleotide.

For example, the SANC may be designed to bind to a receptor found only in a certain cell type, as discussed above. The SANC is also designed to recognize

and selectively bind to target mRNA sequence, which can correspond to a sequence contained within the sequences shown in SEQ ID NOS:1, 3 or 41. The SANC is designed to inactivate target mRNA sequence by either binding thereto and inducing degradation of the mRNA by, for example, RNase I digestion, or inhibiting translation of mRNA target sequence by interfering with the binding of translation-regulating factors or ribosomes, or inclusion of other chemical structures, such as ribozyme sequences or reactive chemical groups which either degrade or chemically modify the target mRNA. SANCs have been shown to be capable of such properties when directed against mRNA targets (see Cohen et al., TIPS, 10:435 (1989) and Weintraub, Sci. American, January (1990), pp.40).

The invention further provides a method of modulating the level of apoptosis in a cell by introducing an antisense nucleotide sequence into the cell, wherein the antisense nucleotide sequence specifically hybridizes to a nucleic acid molecule encoding a Bcl-G, wherein the hybridization reduces or inhibits the expression of the Bcl-G in the cell. The use of anti-sense nucleic acids, including recombinant anti-sense nucleic acids or SANCs, can be advantageously used to inhibit cell death.

Compositions comprising an amount of the antisense-nucleic acid of the invention, effective to reduce expression of Bcl-G polypeptides by entering a cell and binding specifically to mRNA encoding Bcl-G polypeptides so as to prevent translation and an acceptable hydrophobic carrier capable of passing through a cell membrane are also provided herein. Suitable hydrophobic carriers are described, for example, in U.S. Patent Nos. 5,334,761; 4,889,953; 4,897,355, and the

like. The acceptable hydrophobic carrier capable of passing through cell membranes may also comprise a structure which binds to a receptor specific for a selected cell type and is thereby taken up by cells of the selected cell type. For example, the structure can be part of a protein known to bind to a cell-type specific receptor such as a tumor.

Antisense-nucleic acid compositions are useful to inhibit translation of mRNA encoding invention polypeptides. Synthetic oligonucleotides, or other antisense chemical structures are designed to bind to mRNA encoding Bcl-G polypeptides and inhibit translation of mRNA and are useful as compositions to inhibit expression of Bcl-G associated genes in a tissue sample or in a subject.

The invention also provides a method for expression of a Bcl-G polypeptide by culturing cells containing a Bcl-G nucleic acid under conditions suitable for expression of Bcl-G. Thus, there is provided a method for the recombinant production of a Bcl-G of the invention by expressing the nucleic acid sequences encoding Bcl-G in suitable host cells. Recombinant DNA expression systems that are suitable to produce Bcl-G described herein are well-known in the art (see, for example, Ausubel et al., *supra*, 1999). For example, the above-described nucleotide sequences can be incorporated into vectors for further manipulation. As used herein, vector refers to a recombinant DNA or RNA plasmid or virus containing discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof.

The invention also provides vectors containing the Bcl-G nucleic acids of the invention. Suitable expression vectors are well-known in the art and include vectors capable of expressing nucleic acid operatively
5 linked to a regulatory sequence or element such as a promoter region or enhancer region that is capable of regulating expression of such nucleic acid. Appropriate expression vectors include those that are replicable in eukaryotic cells and/or prokaryotic cells and those that
10 remain episomal or those which integrate into the host cell genome.

Promoters or enhancers, depending upon the nature of the regulation, can be constitutive or regulated. The regulatory sequences or regulatory
15 elements are operatively linked to a nucleic acid of the invention such that the physical and functional relationship between the nucleic acid and the regulatory sequence allows transcription of the nucleic acid.

Suitable vectors for expression in prokaryotic
20 or eukaryotic cells are well known to those skilled in the art (see, for example, Ausubel et al., supra, 1999). Vectors useful for expression in eukaryotic cells can include, for example, regulatory elements including the SV40 early promoter, the cytomegalovirus (CMV) promoter,
25 the mouse mammary tumor virus (MMTV) steroid-inducible promoter, Moloney murine leukemia virus (MMLV) promoter, and the like. The vectors of the invention are useful for subcloning and amplifying a Bcl-G nucleic acid molecule and for recombinantly expressing a Bcl-G
30 polypeptide. A vector of the invention can include, for example, viral vectors such as a bacteriophage, a baculovirus or a retrovirus; cosmids or plasmids; and, particularly for cloning large nucleic acid molecules,

bacterial artificial chromosome vectors (BACs) and yeast artificial chromosome vectors (YACs). Such vectors are commercially available, and their uses are well known in the art. One skilled in the art will know or can readily determine an appropriate promoter for expression in a particular host cell.

The invention additionally provides recombinant cells containing Bcl-G nucleic acids of the invention. The recombinant cells are generated by introducing into a host cell a vector containing a Bcl-G nucleic acid molecule. The recombinant cells are transduced, transfected or otherwise genetically modified. Exemplary host cells that can be used to express recombinant Bcl-G molecules include mammalian primary cells; established mammalian cell lines, such as COS, CHO, HeLa, NIH3T3, HEK 293 and PC12 cells; amphibian cells, such as *Xenopus* embryos and oocytes; and other vertebrate cells. Exemplary host cells also include insect cells such as *Drosophila*, yeast cells such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, or *Pichia pastoris*, and prokaryotic cells such as *Escherichia coli*.

In one embodiment, nucleic acids encoding the invention Bcl-G polypeptides can be delivered into mammalian cells, either *in vivo* or *in vitro* using suitable vectors well-known in the art. Suitable vectors for delivering a Bcl-G polypeptide, or a functional fragment thereof to a mammalian cell, include viral vectors such as retroviral vectors, adenovirus, adeno-associated virus, lentivirus, herpesvirus, as well as non-viral vectors such as plasmid vectors. Such vectors are useful for providing therapeutic amounts of a Bcl-G polypeptide (see, for example, U.S. Patent No. 5,399,346, issued March 21, 1995). Delivery of Bcl-G polypeptides

or nucleic acids therapeutically can be particularly useful when targeted to a tumor cell, thereby inducing apoptosis in tumor cells. In addition, where it is desirable to limit or reduce the *in vivo* expression of the invention Bcl-G, the introduction of the antisense strand of the invention nucleic acid is contemplated.

Viral based systems provide the advantage of being able to introduce relatively high levels of the heterologous nucleic acid into a variety of cells. Suitable viral vectors for introducing invention nucleic acid encoding an Bcl-G protein into mammalian cells are well known in the art. These viral vectors include, for example, Herpes simplex virus vectors (Geller et al., Science, 241:1667-1669 (1988)); vaccinia virus vectors (Piccini et al., Meth. Enzymology, 153:545-563 (1987)); cytomegalovirus vectors (Mocarski et al., in Viral Vectors, Y. Gluzman and S.H. Hughes, Eds., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1988, pp. 78-84)); Moloney murine leukemia virus vectors (Danos et al., Proc. Natl. Acad. Sci. USA, 85:6460-6464 (1988); Blaese et al., Science, 270:475-479 (1995); Onodera et al., J. Virol., 72:1769-1774 (1998)); adenovirus vectors (Berkner, Biotechniques, 6:616-626 (1988); Cotten et al., Proc. Natl. Acad. Sci. USA, 89:6094-6098 (1992); Graham et al., Meth. Mol. Biol., 7:109-127 (1991); Li et al., Human Gene Therapy, 4:403-409 (1993); Zabner et al., Nature Genetics, 6:75-83 (1994)); adeno-associated virus vectors (Goldman et al., Human Gene Therapy, 10:2261-2268 (1997); Greelish et al., Nature Med., 5:439-443 (1999); Wang et al., Proc. Natl. Acad. Sci. USA, 96:3906-3910 (1999); Snyder et al., Nature Med., 5:64-70 (1999); Herzog et al., Nature Med., 5:56-63 (1999)); retrovirus vectors (Donahue et al., Nature Med., 4:181-186 (1998); Shackelford et al., Proc. Natl. Acad. Sci. USA,

85:9655-9659 (1988); U.S. Patent Nos. 4,405,712, 4,650,764 and 5,252,479, and WIPO publications WO 92/07573, WO 90/06997, WO 89/05345, WO 92/05266 and WO 92/14829; and lentivirus vectors (Kafri et al., Nature Genetics, 17:314-317 (1997)).

For example, in one embodiment of the present invention, adenovirus-transferrin/polylysine-DNA (TfAdpl-DNA) vector complexes (Wagner et al., Proc. Natl. Acad. Sci., USA, 89:6099-6103 (1992); Curiel et al., Hum. Gene Ther., 3:147-154 (1992); Gao et al., Hum. Gene Ther., 4:14-24 (1993)) are employed to transduce mammalian cells with heterologous Bcl-G nucleic acid. Any of the plasmid expression vectors described herein may be employed in a TfAdpl-DNA complex.

Vectors useful for therapeutic administration of a Bcl-G polypeptide of nucleic acid can contain a regulatory element that provides tissue specific or inducible expression of an operatively linked nucleic acid. One skilled in the art can readily determine an appropriate tissue-specific promotor or enhancer that allows exparssion of a Bcl-G polypeptide or nucleic acid in a desired tissue. Any of a variety of inducible promoters or enhancers can also be included in the vector for regulatable expression of a Bcl-G polypeptide or nucleic acid. Such inducible systems, include, for example, tetracycline inducible system (Gossen & Bizard, Proc. Natl. Acad. Sci. USA, 89:5547-5551 (1992); Gossen et al., Science, 268:1766-1769 (1995); Clontech, Palo Alto, CA); metalothionein promoter induced by heavy metals; insect steroid hormone responsive to ecdysone or related steroids such as muristerone (No et al., Proc. Natl. Acad. Sci. USA, 93:3346-3351 (1996); Yao et al., Nature, 366:476-479 (1993); Invitrogen, Carlsbad, CA);

mouse mammary tumor virus (MMTV) induced by steroids such as glucocorticoid and estrogen (Lee et al., *Nature*, 294:228-232 (1981); and heat shock promoters inducible by temperature changes.

5 An inducible system particularly useful for therapeutic administration utilizes an inducible promoter that can be regulated to deliver a level of therapeutic product in response to a given level of drug administered to an individual and to have little or no expression of
10 the therapeutic product in the absence of the drug. One such system utilizes a Gal4 fusion that is inducible by an antiprogesterone such as mifepristone in a modified adenovirus vector (Burien et al., Proc. Natl. Acad. Sci. USA, 96:355-360 (1999)). Another such inducible system
15 utilizes the drug rapamycin to induce reconstitution of a transcriptional activator containing rapamycin binding domains of FKBP12 and FRAP in an adeno-associated virus vector (Ye et al., Science, 283:88-91 (1999)). It is understood that any combination of an inducible system
20 can be combined in any suitable vector, including those disclosed herein. Such a regulatable inducible system is advantageous because the level of expression of the therapeutic product can be controlled by the amount of drug administered to the individual or, if desired,
25 expression of the therapeutic product can be terminated by stopping administration of the drug.

 The invention additionally provides an isolated anti-Bcl-G antibody having specific reactivity with a Bcl-G. The anti-Bcl-G antibody can be a monoclonal
30 antibody or a polyclonal antibody. The invention further provides cell lines producing monoclonal antibodies having specific reactivity with a Bcl-G.

The invention thus provides antibodies that specifically bind a Bcl-G polypeptide. As used herein, the term "antibody" is used in its broadest sense to include polyclonal and monoclonal antibodies, as well as antigen binding fragments of such antibodies. With regard to an anti-Bcl-G antibody of the invention, the term "antigen" means a native or synthesized Bcl-G polypeptide or fragment thereof. An anti-Bcl-G antibody, or antigen binding fragment of such an antibody, is characterized by having specific binding activity for a Bcl-G polypeptide or a peptide portion thereof of at least about $1 \times 10^5 \text{ M}^{-1}$. Thus, Fab, F(ab')_2 , Fd and Fv fragments of an anti-Bcl-G antibody, which retain specific binding activity for a Bcl-G polypeptide, are included within the definition of an antibody. Specific binding activity of a Bcl-G polypeptide can be readily determined by one skilled in the art, for example, by comparing the binding activity of an anti-Bcl-G antibody to a Bcl-G polypeptide versus a control polypeptide that is not a Bcl-G polypeptide. Methods of preparing polyclonal or monoclonal antibodies are well known to those skilled in the art (see, for example, Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1988)).

In addition, the term "antibody" as used herein includes naturally occurring antibodies as well as non-naturally occurring antibodies, including, for example, single chain antibodies, chimeric, bifunctional and humanized antibodies, as well as antigen-binding fragments thereof. Such non-naturally occurring antibodies can be constructed using solid phase peptide synthesis, can be produced recombinantly or can be obtained, for example, by screening combinatorial libraries consisting of variable heavy chains and

variable light chains as described by Huse et al.
(Science 246:1275-1281 (1989)). These and other methods
of making, for example, chimeric, humanized, CDR-grafted,
single chain, and bifunctional antibodies are well known
5 to those skilled in the art (Winter and Harris, Immunol.
Today 14:243-246 (1993); Ward et al., Nature 341:544-546
(1989) ; Harlow and Lane, *supra*, 1988); Hilyard et al.,
Protein Engineering: A practical approach (IRL Press
1992); Borrabeck, Antibody Engineering, 2d ed. (Oxford
10 University Press 1995)).

Anti-Bcl-G antibodies can be raised using a
Bcl-G immunogen such as an isolated Bcl-G polypeptide
having the amino acid sequence of SEQ ID NOS:2, 4 or 42,
or a fragment thereof, which can be prepared from natural
15 sources or produced recombinantly, or a peptide portion
of the Bcl-G polypeptide. Such peptide portions of a
Bcl-G polypeptide are functional antigenic fragments if
the antigenic peptides can be used to generate a Bcl-G-
specific antibody. A non-immunogenic or weakly
20 immunogenic Bcl-G polypeptide or portion thereof can be
made immunogenic by coupling the hapten to a carrier
molecule such as bovine serum albumin (BSA) or keyhole
limpet hemocyanin (KLH). Various other carrier molecules
and methods for coupling a hapten to a carrier molecule
25 are well known in the art (see, for example, Harlow and
Lane, *supra*, 1988). An immunogenic Bcl-G polypeptide
fragment can also be generated by expressing the peptide
portion as a fusion protein, for example, to glutathione
S transferase (GST), polyHis or the like. Methods for
30 expressing peptide fusions are well known to those
skilled in the art (Ausubel et al., Current Protocols in
Molecular Biology (Supplement 47), John Wiley & Sons, New
York (1999)).

The invention further provides a method for detecting the presence of a human Bcl-G in a sample by contacting a sample with a Bcl-G-specific antibody, and detecting the presence of specific binding of the antibody to the sample, thereby detecting the presence of a human Bcl-G in the sample. Bcl-G specific antibodies can be used in diagnostic methods and systems to detect the level of Bcl-G present in a sample. As used herein, the term "sample" is intended to mean any biological fluid, cell, tissue, organ or portion thereof, that includes or potentially includes Bcl-G nucleic acids or polypeptides. The term includes samples present in an individual as well as samples obtained or derived from the individual. For example, a sample can be a histologic section of a specimen obtained by biopsy, or cells that are placed in or adapted to tissue culture. A sample further can be a subcellular fraction or extract, or a crude or substantially pure nucleic acid or protein preparation.

Bcl-G-specific antibodies can also be used for the immunoaffinity or affinity chromatography purification of the invention Bcl-G. In addition, methods are contemplated herein for detecting the presence of an invention Bcl-G protein in a cell, comprising contacting the cell with an antibody that specifically binds to Bcl-G polypeptides under conditions permitting binding of the antibody to the Bcl-G polypeptides, detecting the presence of the antibody bound to the Bcl-G polypeptide, and thereby detecting the presence of invention polypeptides in a cell. With respect to the detection of such polypeptides, the antibodies can be used for *in vitro* diagnostic or *in vivo* imaging methods.

Immunological procedures useful for *in vitro* detection of target Bcl-G polypeptides in a sample include immunoassays that employ a detectable antibody. Such immunoassays include, for example,

5 immunohistochemistry, immunofluorescence, ELISA assays, radioimmunoassay, FACS analysis, immunoprecipitation, immunoblot analysis, Pandex microfluorimetric assay, agglutination assays, flow cytometry and serum diagnostic assays, which are well known in the art (Harlow and Lane,

10 *supra*, 1988; Harlow and Lane, Using Antibodies: A Laboratory Manual, Cold Spring Harbor Press (1999)).

An antibody can be made detectable by various means well known in the art. For example, a detectable marker can be directly attached to the antibody or

15 indirectly attached using, for example, a secondary agent that recognizes the Bcl-G specific antibody. Useful markers include, for example, radionucleotides, enzymes, binding proteins such as biotin, fluorogens, chromogens and chemiluminescent labels.

As used herein, the terms "label" and "indicating means" in their various grammatical forms refer to single atoms and molecules that are either

20 directly or indirectly involved in the production of a detectable signal. Any label or indicating means can be

25 linked to invention nucleic acid probes, expressed proteins, polypeptide fragments, or antibody molecules. These atoms or molecules can be used alone or in conjunction with additional reagents. Such labels are themselves well-known in clinical diagnostic chemistry.

30 The labeling means can be a fluorescent labeling agent that chemically binds to antibodies or antigens without denaturation to form a fluorochrome

(dye) that is a useful immunofluorescent tracer. A description of immunofluorescent analytic techniques is found in DeLuca, "Immunofluorescence Analysis", in Antibody As a Tool, Marchalonis et al., eds., John Wiley & Sons, Ltd., pp. 189-231 (1982), which is incorporated herein by reference.

In one embodiment, the indicating group is an enzyme, such as horseradish peroxidase (HRP), glucose oxidase, and the like. In another embodiment, radioactive elements are employed labeling agents. The linking of a label to a substrate, i.e., labeling of nucleic acid probes, antibodies, polypeptides, and proteins, is well known in the art. For instance, an invention antibody can be labeled by metabolic incorporation of radiolabeled amino acids provided in the culture medium. See, for example, Galfre et al., Meth. Enzymol., 73:3-46 (1981). Conventional means of protein conjugation or coupling by activated functional groups are particularly applicable. See, for example, Aurameas et al., Scand. J. Immunol., Vol. 8, Suppl. 7:7-23 (1978), Rodwell et al., Biotech., 3:889-894 (1984), and U.S. Patent No. 4,493,795.

In addition to detecting the presence of a Bcl-G polypeptide, invention anti-Bcl-G antibodies are contemplated for use herein to modulate the activity of the Bcl-G polypeptide in living animals, in humans, or in biological tissues or fluids isolated therefrom. The term "modulate" refers to a compound's ability to increase the biological activity by functioning as an agonist or inhibit the biological activity by functioning as an antagonist of an invention Bcl-G polypeptide. Accordingly, compositions comprising a carrier and an amount of an antibody having specificity for Bcl-G

polypeptides effective to block naturally occurring ligands or other Bcl-G-binding proteins from binding to invention Bcl-G polypeptides are contemplated herein. For example, a monoclonal antibody directed to an epitope
5 of an invention Bcl-G polypeptide, including an amino acid sequence set forth in SEQ ID NOS:2, 4 or 42, can be useful for this purpose.

The present invention further provides transgenic non-human mammals that are capable of
10 expressing exogenous nucleic acids encoding Bcl-G polypeptides. As employed herein, the phrase "exogenous nucleic acid" refers to nucleic acid sequence which is not native to the host, or which is present in the host in other than its native environment, for example, as
15 part of a genetically engineered DNA construct. In addition to naturally occurring levels of Bcl-G, a Bcl-G polypeptide of the invention can either be overexpressed or underexpressed in transgenic mammals, for example, underexpressed in a knock-out animal.

Also provided are transgenic non-human mammals capable of expressing nucleic acids encoding Bcl-G polypeptides so mutated as to be incapable of normal activity. Therefore, the transgenic non-human mammals do not express native Bcl-G or have reduced expression of
20 native Bcl-G. The present invention also provides transgenic non-human mammals having a genome comprising antisense nucleic acids complementary to nucleic acids encoding Bcl-G polypeptides, placed so as to be transcribed into antisense mRNA complementary to mRNA
25 encoding Bcl-G polypeptides, which hybridizes to the mRNA and, thereby, reduces the translation thereof. The nucleic acid can additionally comprise an inducible promoter and/or tissue specific regulatory elements, so
30

that expression can be induced, or restricted to specific cell types.

Animal model systems useful for elucidating the physiological and behavioral roles of Bcl-G polypeptides are also provided, and are produced by creating transgenic animals in which the expression of the Bcl-G polypeptide is altered using a variety of techniques. Examples of such techniques include the insertion of normal or mutant versions of nucleic acids encoding an Bcl-G polypeptide by microinjection, retroviral infection or other means well known to those skilled in the art, into appropriate fertilized embryos to produce a transgenic animal, see, for example, Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual (Cold Spring Harbor Laboratory, (1986)). Transgenic animal model systems are useful for *in vivo* screening of compounds for identification of specific ligands, such as agonists or antagonists, which activate or inhibit a biological activity.

Also contemplated herein, is the use of homologous recombination of mutant or normal versions of Bcl-G genes with the native gene locus in transgenic animals, to alter the regulation of expression or the structure of Bcl-G polypeptides by replacing the endogeneous gene with a recombinant or mutated Bcl-G gene. Methods for producing a transgenic non-human mammal including a gene knock-out non-human mammal, are well known to those skilled in the art (see, Capecchi et al., Science 244:1288 (1989); Zimmer et al., Nature 338:150 (1989); Shastry, Experientia, 51:1028-1039 (1995); Shastry, Mol. Cell. Biochem., 181:163-179 (1998); and U.S. Patent No. 5,616,491, issued April 1, 1997, No. 5,750,826, issued May 12, 1998, and No. 5,981,830, issued

November 9, 1999).

Invention nucleic acids, oligonucleotides, including antisense, vectors containing invention nucleic acids, transformed host cells, polypeptides and combinations thereof, as well as antibodies of the present invention, can be used to screen compounds to determine whether a compound functions as a potential agonist or antagonist of invention polypeptides. These screening assays provide information regarding the function and activity of invention polypeptides, which can lead to the identification and design of compounds that are capable of specific interaction with one or more types of polypeptides, peptides or proteins.

Thus, the invention provides methods for identifying compounds which bind to Bcl-G polypeptides. The invention proteins can be employed in a competitive binding assay. Such an assay can accommodate the rapid screening of a large number of compounds to determine which compounds, if any, are capable of binding to Bcl-G polypeptides. Subsequently, more detailed assays can be carried out with those compounds found to bind, to further determine whether such compounds act as modulators, agonists or antagonists of invention Bcl-G polypeptides. Compounds that bind to and/or modulate invention Bcl-G polypeptides can be used to treat a variety of pathologies mediated by invention Bcl-G polypeptides.

Various binding assays to identify cellular proteins that interact with protein binding domains are known in the art and include, for example, yeast two-hybrid screening assays (see, for example, U.S. Patent Nos. 5,283,173, 5,468,614 and 5,667,973; Ausubel

et al., *supra*, 1999; Luban et al., Curr. Opin. Biotechnol. 6:59-64 (1995)) and affinity column chromatography methods using cellular extracts. By synthesizing or expressing polypeptide fragments
5 containing various Bcl-G sequences or deletions, the Bcl-G binding interface can be readily identified.

In another embodiment of the invention, there is provided a bioassay for identifying compounds which modulate the activity of invention Bcl-G polypeptides.
10 According to this method, invention polypeptides are contacted with an "unknown" or test substance, for example, in the presence of a reporter gene construct responsive to a Bcl-G signaling pathway, the activity of the polypeptide is monitored subsequent to the contact
15 with the "unknown" or test substance, and those substances which cause the reporter gene construct to be expressed are identified as functional ligands for Bcl-G polypeptides. Such reporter gene assays and systems are well known to those skilled in the art (Ausubel et al.,
20 *supra*, 1999). In addition, a reporter gene construct can be generated using the promoter region of Bcl-G and screened for compounds that increase or decrease Bcl-G gene promoter activity. Such compounds can also be used to alter Bcl-G expression.

25 In accordance with another embodiment of the present invention, transformed host cells that recombinantly express invention polypeptides can be contacted with a test compound, and the modulating effect(s) thereof can then be evaluated by comparing the
30 Bcl-G-mediated response, for example, via reporter gene expression in the presence and absence of test compound, or by comparing the response of test cells or control cells, to the presence of the compound.

As used herein, a compound or a signal that "modulates the activity" of invention polypeptides refers to a compound or a signal that alters the activity of Bcl-G polypeptides so that the activity of the invention polypeptide is different in the presence of the compound or signal than in the absence of the compound or signal. In particular, such compounds or signals include agonists and antagonists. An agonist encompasses a compound or a signal that activates Bcl-G protein expression or biological activity. Alternatively, an antagonist includes a compound or signal that interferes with Bcl-G expression or biological activity. Typically, the effect of an antagonist is observed as a blocking of agonist-induced protein activation. Antagonists include competitive and non-competitive antagonists.

Assays to identify compounds that modulate Bcl-G polypeptide expression can involve detecting a change in Bcl-G polypeptide abundance in response to contacting the cell with a compound that modulates Bcl-G activity. Assays for detecting changes in polypeptide expression include, for example, immunoassays with Bcl-G-specific Bcl-G antibodies, such as immunoblotting, immunofluorescence, immunohistochemistry and immunoprecipitation assays, as described above.

As understood by those of skill in the art, assay methods for identifying compounds that modulate Bcl-G activity generally require comparison to a control. One type of a "control" is a cell or culture that is treated substantially the same as the test cell or test culture exposed to the compound, with the distinction that the "control" cell or culture is not exposed to the compound. Another type of "control" cell or culture can be a cell or culture that is identical to the test cells,

with the exception that the "control" cells or culture do not express a Bcl-G polypeptide. Accordingly, the response of the transfected cell to a compound is compared to the response, or lack thereof, of the "control" cell or culture to the same compound under the same reaction conditions.

Methods for producing pluralities of compounds to use in screening for compounds that modulate the activity of a Bcl-G polypeptide, including chemical or biological molecules such as simple or complex organic molecules, metal-containing compounds, carbohydrates, peptides, proteins, peptidomimetics, glycoproteins, lipoproteins, nucleic acids, antibodies, and the like, are well known in the art and are described, for example, in Huse, U.S. Patent No. 5,264,563; Francis et al., Curr. Opin. Chem. Biol. 2:422-428 (1998); Tietze et al., Curr. Biol., 2:363-371 (1998); Sofia, Mol. Divers. 3:75-94 (1998); Eichler et al., Med. Res. Rev. 15:481-496 (1995); and the like. Libraries containing large numbers of natural and synthetic compounds also can be obtained from commercial sources. Combinatorial libraries of molecules can be prepared using well known combinatorial chemistry methods (Gordon et al., J. Med. Chem. 37: 1233-1251 (1994); Gordon et al., J. Med. Chem. 37: 1385-1401 (1994); Gordon et al., Acc. Chem. Res. 29:144-154 (1996); Wilson and Czarnik, eds., Combinatorial Chemistry: Synthesis and Application, John Wiley & Sons, New York (1997)).

Compounds that modulate Bcl-G activity can be screened by the methods disclosed herein to identify compounds that modulate any biological activity or function of Bcl-G. For example, compounds can be identified that alter the interaction of Bcl-G with Bcl-2

family members. Additionally, compounds can be identified that modulate ion channel activity associated with Bcl-G. The formation of ion channels by Bcl-2 family members is one mechanism of inducing apoptosis in cells (Reed, *supra*, 1998). Therefore, compounds that modulate ion channel activity of Bcl-G can be used to alter apoptosis, thereby increasing or decreasing apoptotic activity of Bcl-G.

Another assay for screening of compounds that modulates the activity of Bcl-G is based on altering the phenotype of yeast by expressing Bcl-G. For example, expression of Bax in yeast confers a lethal phenotype (Matsuyama et al., Mol. Cell. 1:327-336 (1998)). A yeast that expresses Bcl-G can have a similar phenotype as Bax since the biological activity of Bcl-G is similar to Bax (Example III). Accordingly, a yeast strain expressing Bcl-G that confers a lethal phenotype can be screened for compounds that prevent cell death. In one embodiment, expression of Bcl-G can be inducible (Tao et al., J. Biol. Chem. 273:23704-23708 (1998), and the compounds can be screened when Bcl-G expression is induced. Bcl-G can also be co-expressed in yeast with other Bcl-2 family members having anti-apoptotic activity such as Bcl-2 or Bcl-X_L. For example, co-expression of Bax with Bcl-2 or Bcl-X_L suppressed the lethal activity of Bax (Tao et al., *supra*, 1998). Similarly, co-expression of Bcl-G with an anti-apoptotic Bcl-2 family member such as Bcl-2 or Bcl-X_L can be used to screen for compounds that antagonize the activity of the anti-apoptotic Bcl-2 family members and restore a lethal phenotype. Such compounds can function to inhibit binding of Bcl-G to anti-apoptotic Bcl-2 family members such as Bcl-2 or Bcl-X_L.

In yet another embodiment of the present invention, the activation of Bcl-G polypeptides can be modulated by contacting the polypeptides with an effective amount of at least one compound identified by the assays described herein. The invention also provides a method of identifying an effective agent that alters the association of a Bcl-G with a Bcl-G associated polypeptide (BAP). The method includes the steps of contacting the Bcl-G and the BAP polypeptide, under conditions that allow said Bcl-G and BAP polypeptide to associate, with a compound; and detecting the altered association of the Bcl-G and BAP polypeptide, thereby identifying a compound that is an effective agent for altering the association of Bcl-G with BAP. The compound can be, for example, a drug or polypeptide. A BAP can be, for example, Bcl-2 family member such as Bcl-2 or Bcl-X_L.

As disclosed herein, Bcl-G is a new member of the Bcl-2 family that has pro-apoptotic activity (see Example III). Therefore, modulation of Bcl-G activity can be advantageously used to modulate the level of apoptosis in a cell. For example, increasing the activity of Bcl-G can be used to promote apoptosis in a cell. Bcl-G activity can be increased, for example, by increasing the level of a Bcl-G polypeptide or functional fragment thereof. Increased levels of a Bcl-G polypeptide can be accomplished, for example, by delivering to a cell a nucleic acid encoding Bcl-G and expressing a Bcl-G polypeptide recombinantly or by delivering a Bcl-G polypeptide or functional fragment thereof directly to a target by the methods disclosed herein. Additionally, Bcl-G activity can be increased by using a modulatory agent that functions as an agonist. Promoting apoptosis by increasing Bcl-G activity or

expression is useful, for example, in therapeutic applications such as the treatment of cancer.

As disclosed herein, decreases or loss of Bcl-G is associated with approximately 50% of prostate cancers, approximately 30% of ovarian cancers and approximately 30 % of leukemias. Bcl-G can function as a tumor suppressor. Therefore, methods of administering a Bcl-G polypeptide either directly or using an encoded nucleic acid can be used to treat a cancer. Furthermore, many chemotherapeutic agents function through increasing apoptosis. Therefore, the invention additionally provides a method to enhance a chemotherapy by increasing Bcl-G activity or expression. Administering Bcl-G can thus be used to enhance the effect of standard chemotherapeutic agents.

Alternatively, modulation of Bcl-G activity can be advantageously used to decrease Bcl-G activity to decrease apoptosis. For example, Bcl-G activity or expression can be decreased by administering an anti-sense Bcl-G nucleic acid. In addition, an antagonist of Bcl-G activity can be identified by the methods disclosed herein and used to decrease Bcl-G activity. Decreasing Bcl-G activity can be used to inhibit apoptosis. Inhibiting apoptosis can be useful, for example, to treat disease ischemic. For example, decreasing Bcl-G activity with anti-sense nucleic acids or small molecule compounds can be used to treat stroke, heart attack, autoimmunity, trauma, neuron cell death, and inflammatory diseases, including Crohn's disease. For example, Bcl-G was identified in Crohn's disease patients (see Example I).

The invention further provides a method for modulating an activity mediated by a Bcl-G polypeptide by

contacting the Bcl-G polypeptide with an effective,
modulating amount of an agent that modulates Bcl-G
activity. The Bcl-G activity can be, for example,
apoptosis-inducing activity, binding to Bcl-2, or tumor
5 suppressor activity. The invention additionally provides
a method of modulating the level of apoptosis in a cell.
The method includes the steps of introducing a nucleic
acid molecule encoding a Bcl-G into the cell; and
expressing the Bcl-G in the cell, wherein the expression
10 of the Bcl-G modulates apoptosis in the cell.

The invention further provides a method of
modulating the level of apoptosis in a cell by contacting
the cell with a compound that effectively alters the
association of Bcl-G with a Bcl-G-associated-protein in
15 the cell, or that effectively alters the activity of a
Bcl-G in the cell. Additionally provided by the
invention is a method of modulating interactions between
Bcl-G and Bcl-2 by contacting a Bcl-G polypeptide with
the agent that inhibits or alters interactions between
20 Bcl-G and Bcl-2.

As disclosed herein, Bcl-G is located on
chromosome 12 in a region deleted in various cancers,
including leukemia, prostate and ovarian cancer (Example
IV). Therefore, methods using Bcl-G nucleic acids or
25 antibodies can be used as a diagnostic for predisposition
or progression of cancer, for example, leukemia, prostate
or ovarian cancer. Changes in Bcl-G expression or
activity can be correlated with patient survival or
response to therapy, and a correlation can be used to
30 monitor cancer progression or response to therapy.

The invention further provides a method of
diagnosing a pathology characterized by an increased or

decreased level of a Bcl-G in a subject. The method includes the steps of (a) obtaining a test sample from the subject; (b) contacting the sample with an agent that can bind the Bcl-G under suitable conditions, wherein the conditions allow specific binding of the agent to the Bcl-G; and (c) comparing the amount of the specific binding in the test sample with the amount of specific binding in a control sample, wherein an increased or decreased amount of the specific binding in the test sample as compared to the control sample is diagnostic of a pathology. The agent can be, for example, an anti-Bcl-G antibody, a Bcl-G-associated-protein (BAP), or a Bcl-G nucleic acid.

The invention also provides a method of diagnosing cancer or monitoring cancer therapy by contacting a test sample from a patient with a Bcl-G-specific antibody. The invention additionally provides a method of assessing prognosis of patients with cancer comprising contacting a test sample from a patient with a Bcl-G-specific antibody.

The invention additionally provides a method of diagnosing cancer or monitoring cancer therapy by contacting a test sample from a patient with a Bcl-G oligonucleotide. The invention further provides a method of assessing prognosis of patients with cancer by contacting a test sample from a patient with a Bcl-G oligonucleotide.

The methods of the invention for diagnosing cancer or monitoring cancer therapy using a Bcl-G-specific antibody or Bcl-G oligonucleotide or nucleic acid can be used, for example, to segregate patients into a high risk group or a low risk group for predicting risk

of metastasis or risk of failure to respond to therapy. Therefore, the methods of the invention can be advantageously used to determine the risk of metastasis in a cancer patient or as a prognostic indicator of survival in a cancer patient. One of ordinary skill in the art would appreciate that the prognostic indicators of survival for cancer patients suffering from stage I cancer can be different from those for cancer patients suffering from stage IV cancer. For example, prognosis for stage I cancer patients can be oriented toward the likelihood of continued growth and/or metastasis of the cancer, whereas prognosis for stage IV cancer patients can be oriented toward the likely effectiveness of therapeutic methods for treating the cancer. Accordingly, the methods of the invention directed to measuring the level of or determining the presence of a Bcl-G polypeptide or encoding nucleic acid can be used advantageously as a prognostic indicator for the presence or progression of a cancer or response to therapy.

In accordance with another embodiment of the present invention, there are provided diagnostic systems, preferably in kit form, comprising at least one invention nucleic acid or antibody in a suitable packaging material. The diagnostic kits containing nucleic acids are derived from the Bcl-G-encoding nucleic acids described herein. In one embodiment, for example, the diagnostic nucleic acids are derived from any of SEQ ID NOS:1, 3 or 41 and can be oligonucleotides of the invention. Invention diagnostic systems are useful for assaying for the presence or absence of nucleic acid encoding Bcl-G in either genomic DNA or mRNA.

A suitable diagnostic system includes at least one invention nucleic acid or antibody, as a separately

packaged chemical reagent(s) in an amount sufficient for at least one assay. For a diagnostic kit containing nucleic acid of the invention, the kit will generally contain two or more nucleic acids. When the diagnostic
5 kit is to be used in PCR, the kit will contain at least two oligonucleotides that can serve as primers for PCR. Those of skill in the art can readily incorporate invention nucleic probes and/or primers or invention
10 antibodies into kit form in combination with appropriate buffers and solutions for the practice of the invention methods as described herein. A kit containing a Bcl-G antibody can contain a reaction cocktail that provides the proper conditions for performing an assay, for
15 example, an ELISA or other immunoassay, for determining the level of expression of a Bcl-G polypeptide in a sample, and can contain control samples that contain known amounts of a Bcl-G polypeptide and, if desired, a second antibody specific for the anti-Bcl-G antibody.

The contents of the kit of the invention, for
20 example, Bcl-G nucleic acids or antibodies, are contained in packaging material, preferably to provide a sterile, contaminant-free environment. In addition, the packaging material contains instructions indicating how the
25 materials within the kit can be employed both to detect the presence or absence of a particular Bcl-G sequence or Bcl-G polypeptide or to diagnose the presence of, or a predisposition for a condition associated with the
30 presence or absence of Bcl-G such as cancer. The instructions for use typically include a tangible expression describing the reagent concentration or at least one assay method parameter, such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent/sample admixtures, temperature, buffer conditions, and the like.

It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also provided within the definition of the invention provided herein.

5 Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Molecular Cloning of Bcl-G

10 This example describes the cloning of Bcl-G, a homologue of Bcl-2.

To clone the full length Bcl-G gene, oligonucleotide primers were designed based a short EST (GenBank Accession No. AW000827) from colonic mucosa of 3 patients with Crohn's disease found by searching a

15 database for sequences similar to the BH2 and BH3 domains of Bcl-2 family proteins. The primers used were Primer 1 (5'-GTACTTGGTGCCAAAGCCCAGG-3'; SEQ ID NO:7) and Primer 2 (5'-GACATGATGTCTGGTGTAGTAGGCGAGG-3'; SEQ DI NO:8). The full length Bcl-G cDNA was cloned using SMARTTMRACE cDNA

20 Amplification Kit (Clontech; Palo Alto CA) from human placental total RNA (Clontech) as template. The 5'-RACE products were sequenced with an automated sequencer.

Briefly, for cloning of Bcl-G cDNAs, TBLAST searches of the public databases using human Bcl-2 as a

25 query sequence revealed a short EST (GenBank AW000827) from colonic mucosa of 3 patients with Crohn's disease which contains an open reading frame (ORF) encoding sequences similar to the BH2 domain of Bcl-2 family proteins. An oligonucleotide primer

30 (5'-GTACTTGGTGCCAAAGCCCAGG-3'; SEQ ID NO:7) was designed complementary to the EST sequence and used for 5'-RACE,

employing the SMART™ RACE cDNA Amplification Kit (Clontech; Palo Alto CA) and human placental total RNA as template. The 5'-RACE products were subcloned into pCR2.1-TOPO vector using the TOPO™ TA Cloning kit (Invitrogen; Carlsbad CA), and their DNA sequence determined, revealing a complete open reading frame (ORF), with start codon within a favorable Kozak sequence context, preceded by a 5' -untranslated region (UTR) containing stop codons in all three reading-frames. (submitted to Genbank). Two additional EST clones, AI478889 and AI240211, were identified by BLAST searches, corresponding to overlapping partial Bcl-G cDNAs which contained the 3' -UTR.

A short EST was identified during searches of the public databases, which when conceptually translated revealed a polypeptide sequence with similarity to the BH2 domain of Bcl-2 family proteins. Full-length cDNAs were obtained, revealing two potential transcripts containing open reading frames (ORF) for proteins of 327 and 252 amino-acids, respectively, which were termed Bcl-G_L and Bcl-G_S (Figure 5A). The predicted Bcl-G_L and Bcl-G_S proteins are identical for the first 226 amino acids, then diverge thereafter. Comparison of the predicted amino acid sequences of Bcl-G_L and Bcl-G_S with Bcl-2 family proteins revealed the presence of a candidate BH3 domain (SEQ ID NO:9) in both Bcl-G_L and Bcl-G_S (Figure 5A,B), and the presence of a BH2 domain (SEQ ID NO:18) in Bcl-G_L but not in Bcl-G_S (Figure 5A, C).

Invention Bcl-G was found to exist in two forms, a long form, designated Bcl-G_L, and a shorter form, designated Bcl-G_S. The nucleotide sequence of Bcl-G_L is shown in Figure 1 (SEQ ID NO:1). The nucleotide sequence of the coding region of Bcl-G_L cDNA and the encoded amino

acid sequence (SEQ ID NO:3) are shown in Figure 2. Bcl-G_L was initially identified to contain a core BH3 domain (216LKYSGDQLE²²⁴; SEQ ID NO:5) and a core BH2 domain (307PWIQQHGGWE³¹⁶; SEQ ID NO:6).

5 The shorter form of Bcl-G, Bcl-G_S, is an apparent alternative splicing product of Bcl-G mRNA. The nucleotide sequence of Bcl-G_S is shown in Figure 3 (SEQ ID NO:3). The nucleotide sequence of the coding region of Bcl-G_S cDNA and the encoded amino acid sequence (SEQ ID
10 NO:4) are shown in Figure 2. Bcl-G_S contains only the BH3 domain (216LKYSGDQLE²²⁴).

 These results demonstrate that a new member of the Bcl-2 family, Bcl-G, is expressed in human placenta and in the colonic mucosa of patients with Crohn's
15 disease. Bcl-G exists in two forms, a long form, designated Bcl-G_L, which contains a BH2 and BH3 domain, and a short form, designated Bcl-G_S, which contains only a BH3 domain.

EXAMPLE II

20 Mapping of Bcl-G to Chromosome 12p12.3

 This experiment describes chromosomal mapping of human Bcl-G.

 To map the chromosomal location of Bcl-G, a search of the GenBank database was performed using BLAST
25 (Altschul et al., J. Mol. Biol. 215:403-410 (1990); Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). A 190858 bp human 12p12 BAC chromosome sequence RPCI11-267J23 (GenBank accession no. AC007537) was found to contain the full length Bcl-G gene. The BAC also
30 contains the LRP6 gene (exon 1 starts at 89963 bp). A

600 kb region between 12p12.3 to 12p13.1, flanked by D12S358 and ETV6/exon8, was previously defined to be frequently deleted in childhood acute lymphoblastic leukemia (ALL) and other solid tumor cells (Baens et al., (1999) Genomics 56:40-50 (1999); Hatta et al., Br. J. Cancer 75:1256-1262 (1997); Kibel et al., Cancer Res. 58:5652-5655 (1998); Baccichet et al., Br. J. Haematol. 99:107-114 (1997); Aissani et al., Leuk. Lymphoma 34:231-239). The loss of the region containing Bcl-G occurs in approximately 50% of prostate cancers, 30% of ovarian cancers, and 30% of leukemias.

The LRP6 gene is located in the region between 12p12.3 to 12p13.1. Using LRP6 as a marker for orientation, Bcl-G was located in this region. Exon 1 of Bcl-G starts at 40674 bp in the BAC and was deduced from novel DNA sequence data obtained from 5' RACE-based amplification of the full-length Bcl-G cDNA. The genomic structure of the Bcl-G gene is shown in Figure 5. Bcl-G_L has 6 exons, with the first codon non-coding, spreading across a 30 kb region in chromosome 12. Bcl-G_S also has 6 exons, but a 153 bps sequence is inserted in front of exon 5 and contains a stop codon. The BH3 domain is located in exon 4 of both Bcl-G_L and Bcl-G_S. The BH2 domain is located in exon 5 of Bcl-G_L. The Bcl-G_L and Bcl-G_S cDNAs can be accounted for by an alternative mRNA splicing mechanism in which different splice acceptor sites associated with exon 5 are employed, resulting in a change in the distal reading-frame (Figure 5D).

The chromosomal mapping of Bcl-G to chromosome 12p12.3 is shown in Figure 6. Bcl-G is located in a 600 kb region that has been previously determined to be frequently deleted in childhood ALL and other solid tumors (Baens et al., *supra*, 1999). Therefore, Bcl-G is

located in a region deleted in ALL and can function as a tumor suppressor or as a marker for tumor suppressor activity.

Example III

Expression of Bcl-G

This example describes the expression of Bcl-G.

For generation of plasmids, cDNAs containing the ORFs of Bcl-G_L and Bcl-G_S without additional flanking sequences were generated by PCR using human placental cDNA as a template and the following primers:

5'-GGCTCGAGCGATGTGTAGCACCAGTGGGTGTGACC-3' (SEQ ID NO:27), sense for both Bcl-G_L and Bcl-G_S;

5'-CCAAGCTTTAAGTCTACTTCTTCATGTGATATCCC-3' (SEQ ID NO:28), antisense for Bcl-G_L; and

5'-CCAAGCTTTAAATGCAGGCCATCAAACC-3' (SEQ ID NO:29), antisense for Bcl-G_S. The resulting PCR products were digested with restriction endonucleases and subcloned into the Xho I and Hind III sites of pEGFP-C1 (Clontech). A mutant of Bcl-G_S lacking the BH3 domain was created by a two-step PCR method, using the following primers:

primer1, 5'- GGCTCGAGCGATGTGTAGCACCAGTGGGTGTGACC-3' (SEQ ID NO:30); primer2,

5'-CCGGATCCGGCTAGTATTTGTTCTTCTTCATCTTTC-3' (SEQ ID NO:31); primer3, 5'-CCGGATCCGACACTGCCTTCATCCCCATTCCC-3' (SEQ ID NO:32); and primer4,

5'-CCAAGCTTTAAATGCAGGCCATCAAACC-3' (SEQ ID NO:33). The resulting PCR product was digested with XhoI / BamHI or with BamHI / HindIII respectively, and ligated into pEGFP-C1. Site-directed mutagenesis of Bcl-G_S was performed to generate a L216E substitution mutation using the QuikChange™ Site-Directed Mutagenesis Kit (Stratagene) following manufacturer's procedure, with

PEGFP-C1/Bcl-G_s plasmid as DNA template, and the mutagenic primers: 5'-GCCAAAATTGTTGAGCTGGAGAAATATTCAGGAGATCAGTTGG-3' (SEQ ID NO:34) and
5'-CCAACTGATCTCCTGAATATTTCTCCAGCTCAACAATTTTGGC-3' (SEQ ID
5 NO:35).

For measurements of Bcl-G mRNAs, Bcl-G mRNAs were detected by either Northern blotting or Reverse-Transcriptase-Polymerase Chain Reaction (RT-PCR). For RT-PCR, multiple-tissue cDNA panels (Clontech) containing
10 first-strand cDNA generated from 16 different tissues were employed. PCR was performed according to the manufacturer's protocol with following primers: (a) 5' primer for both Bcl-G_s and Bcl-G_L, corresponding to exon 3, 5'-CTGAGGGTCTCTCCTCCAGCTCCAAGG-3' (SEQ ID NO:36); (b)
15 3' primer for Bcl-G_L, corresponding to exon 5, 5'-GGCCGTGACGTCTATTACAAGGGCAGCC-3' (SEQ ID NO:37); and 3' primer for Bcl-G_s, corresponding to an alternatively spliced segment of exon 5,
5'-CAAGGGAATGGGGATGAAGGCAGTGTC-3' (SEQ ID NO:38). Human
20 G3PDH expression was examined by PCR with the following primers: 5'-TGAAGGTCGGAGTCAACGGATTGTT-3' (SEQ ID NO:39) (sense); and 5'-CATGTGGGCCATGAGGTCCACCAC-3' (SEQ ID NO:40) (antisense).

For tissue-specific expression of Bcl-G_L and
25 Bcl-G_s mRNAs, Northern blotting demonstrated the presence of ~2 kbp Bcl-G transcripts in several normal human tissues, but failed to resolve the mRNAs encoding Bcl-G_L and Bcl-G_s. RT-PCR assays were therefore designed using primers specific for Bcl-G_L and Bcl-G_s sequences
30 associated with exon 5. Bcl-G_L mRNA was clearly detected in lung, pancreas, prostate and testis, with lower levels present in some other tissues (Figure 7). In contrast, Bcl-G_s mRNA was uniquely expressed in testis. RT-PCR

amplification of a control mRNA, G3PDH, demonstrated loading of nearly equivalent amounts of mRNA from each tissue. The amplified bands corresponding to Bcl-G_L and Bcl-G_S were excised and sequenced, confirming the validity of the RT-PCR strategy.

EXAMPLE IV

Induction of Cell Death by Bcl-G_S

This experiment describes the induction of cell death by Bcl-G_S in transfected PC-3 cells.

For cell culture, transfections, and apoptosis assay, 293T and Cos-7 cells were cultured in DMEM high glucose media (Irvine Scientific, Santa Ana, CA) containing 10% fetal bovine serum (FBS). PC-3 cells were cultured with RPMI 1640 media containing 10% FBS. Transfection of cells was performed using SuperFect (Qiagen, Chatsworth, CA). Both floating and adherent cells (after trypsinization) were collected 24 hrs after transfection, fixed, and stained using 4',6-diamidine-2'-phenylindole dihydrochloride (DAPI) for assessing apoptosis based on nuclear fragmentation and chromatin condensation (Xu & Reed, Mol. Cell, 1:337-346 (1998); Zhang et al., Proc. Natl. Acad. Sci. (USA), 97:2597-2602 (2000)).

To characterize a biological function of Bcl-G, a Bcl-G_S construct was generated by cloning Bcl-G_S cDNA into pcDNA3.1/Myc/His expression vector (Invitrogen; Carlsbad CA) at the Xho I / Hind III sites. The authenticity of the construct was confirmed by DNA sequencing.

For transfection experiments, various vectors were transfected into PC-3 cells: control vector pcDNA3.1/Myc/His; pcDNA3.1/Myc/His/Bcl-G_s expressing Bcl-G_s; pRC/CMV/Bcl-2 expressing Bcl-2; and pRC/CMV/Bax expressing Bax. The vectors were transfected as follows: pcDNA3.1/Myc/His alone; pcDNA3.1/Myc/His/Bcl-G_s alone; pcDNA3.1/Myc/His/Bcl-G_s + pRC/CMV/Bcl-2; pRC/CMV/Bax alone; or pRC/CMV/Bax + pRC/CMV/Bcl-2. One µg of each vector was combined with 0.2 µg pEGFP-N2 (Clontech), and the vectors were transiently transfected into PC-3 prostate cancer cells using SuperFect reagent (QIAGEN; Valencia CA), following the instructions of the manufacturer. At 24 hours after transfection, cells were examined under a fluorescent microscope. About 100 green fluorescent protein (GFP) positive (green color) cells were counted for each transfection. Cells that were detached with membrane blebbing and/or apoptotic bodies were recorded as dead cells. Results were averaged from three separate transfections.

As shown in Figure 8, Bcl-G_s induces cell death in PC-3 cells (compare "control" to "Bcl-G_s"). The induction of cell death by Bcl-G_s was similar to Bax, which was used as a positive control based on its known pro-apoptotic activity (compare "Bcl-G_s" to "Bax"). The induction of cell death by Bcl-G_s was completely inhibited when co-transfected with the anti-apoptotic Bcl-2 (see "Bcl-G_s + Bcl-2"). The inhibition of Bcl-G_s-induced cell death by Bcl-2 was similar to that seen with Bax (see "Bax + Bcl-2").

To assess the effects of Bcl-G_L and further assess the effects of Bcl-G_s on apoptosis, various cell lines, including Cos7, HEK293T, and PC3, were transiently transfected with plasmids encoding Bcl-G_L or Bcl-G_s. For

most experiments, Bcl-G_L and Bcl-G_S were expressed as GFP-fusions so that successfully transfected cells could be conveniently identified (Figure 9A), but similar results were obtained when Flag-epitope tags were employed instead. Over-expression of the shorter Bcl-G_S protein reproducibly induced striking increases in the percentage of cells undergoing apoptosis, as determined by DAPI staining (Figure 9) and other methods. In contrast, Bcl-G_L was more variable and less efficient at inducing apoptosis in these transient transfection assays. Immunoblot analysis of lysates from transfected cells demonstrated that the less potent effects of Bcl-G_L could not be accounted for by lower levels of protein production (Figure 9A). Indeed, Bcl-G_L protein accumulated to levels ~10-fold higher in cells compared to Bcl-G_S, suggesting that Bcl-G_S is a far more potent apoptosis-inducer. Analysis of the same blots with an anti-tubulin antibody confirm loading of essentially equivalent amounts of total protein for each sample, thus validating the results. In additional transfection experiments, Bcl-G_L failed to demonstrate cytoprotective activity in side by side comparisons with Bcl-2 and Bcl-X_L.

EXAMPLE V

The BH3 Domain of Bcl-G_S is Required for its Pro-apoptotic Activity

The Bcl-G_S protein contains a BH3 domain, but lacks other regions of homology with Bcl-2 family proteins. Structural studies indicate that BH3 domains represent amphipathic α -helices, in which the hydrophobic surface of the α -helices of apoptosis-inducing BH3 peptides bind to a pocket on survival proteins such as Bcl-X_L (Sattler et al., Science, 275:983-986 (1997)).

Therefore, the apoptosis-inducing activity of the wild-type Bcl-G_s protein was compared with mutants lacking the BH3 domain (Δ BH3) or in which leucine 216 within the BH3 domain of Bcl-G_s was chosen for mutation to charged glutamic acid, based on comparisons with previously described BH3 mutagenesis experiments demonstrating a critical requirement for the analogous leucine in other pro-apoptotic Bcl-2 family proteins (Wang et al., Mol. Cell. Biol., 18:6083-6089 (1998); Kelekar et al., Mol. Cell. Biol., 17:7040-7046 (1997)).

Wild-type Bcl-G_s potentially induced apoptosis when overexpressed in Cos-7, PC3, HEK293T and other cell lines, whereas Bcl-G_s (Δ BH3) and Bcl-G_s (L216E) did not (see Figure 3B). Immunoblot analysis confirmed production of the Bcl-G_s (Δ BH3) and Bcl-G_s (L216E) proteins at levels exceeding the amounts of wild-type Bcl-G_s protein. Therefore, the BH3 domain of Bcl-G_s is critical for its pro-apoptotic activity.

EXAMPLE VI

Bcl-G_s Associates with Bcl-X_L in a BH3-dependent Manner

The pro-apoptotic activity of "BH3-only" members of the Bcl-2 family depends on their ability to dimerize with and suppress the activity of survival proteins such as Bcl-X_L (reviewed in Kelekar & Thompson, Trends Cell Biol., 8:324-330 (1998)). It was, therefore, determined whether Bcl-G_L and Bcl-G_s are capable of associating with other Bcl-2 family proteins by co-immunoprecipitation assays.

For co-immunoprecipitations and immunoblotting, immunoblotting was performed as described previously (Xu and Reed, supra., (1998); Zhang et al., supra., (2000)).

For co-immunoprecipitations, cells were cultured in 50 mM benzocarbonyl Valine Alanine Aspartate fluoromethylketone (zVAD-fmk) to prevent apoptosis. Cells were suspended in lysis buffer (50 mM Tris-HCl, pH7.4; 150 mM NaCl; 20 mM EDTA; 50 mM NaF; 0.5% NP-40; 0.1 mM Na_3VO_4 ; 20 $\mu\text{g}/\text{ml}$ Leupeptin; 20 $\mu\text{g}/\text{ml}$ Aprotinin; 1 mM dithiothreitol (DTT); and 1 mM phenylmethylsulfonylfluoride (PMSF). Lysates (0.2 ml diluted into 1 ml final volume of lysis buffer) were cleared by incubation with 15 μl of protein G-Sepharose 4B (Zymed; South San Francisco CA) and then incubated with 15 μl of polyclonal anti-GFP antibody (Santa Cruz; Santa Cruz CA) and 15 μl of protein G at 4°C overnight. Beads were then washed 4 times with 1.5 mls lysis buffer before boiling in Laemmli sample buffer and performing SDS-PAGE/immunoblotting.

Bcl- G_s association with the survival proteins. Bcl- X_L and Bcl-2 was readily detected by co-immunoprecipitation using lysates from transiently transfected cells, whereas no association with pro-apoptotic proteins Bax, Bak, Bid or Bad was observed (Figure 10A). Interaction of Bcl- G_s with Bcl-2 and Bcl- X_L , but not with Bax or Bak, was also confirmed by yeast two-hybrid assays. In contrast, association of the longer Bcl- G_L protein with Bcl-2 or Bcl- X_L was not easily detected by co-immunoprecipitation assays (Figure 10A). With much longer x-ray film exposure times, however, small amounts of Bcl- X_L were observed in association with Bcl- G_L immunocomplexes, suggesting either low affinity binding of Bcl- G_L to Bcl- X_L or implying that only a small portion of total Bcl- G_L proteins are competent to bind Bcl- X_L . The interaction of Bcl- G_s with Bcl- X_L was BH3-dependent, as determined by comparisons of wild-type Bcl- G_s with the Bcl- G_s (ΔBH3) and Bcl- G_s (L216E) proteins (Figures 10B, C). Thus, the pro-apoptotic activity of

Bcl-G_s correlates with its ability to bind Bcl-X_L.

EXAMPLE VII

Bcl-G_s is Associated with Cytosolic Organelles

Many Bcl-2 family proteins, such as Bcl-2,
5 Bcl-X_L, and Bak, contain a hydrophobic stretch of amino-
acids near their carboxyl-terminus that anchors them in
intracellular membranes of mitochondria, endoplasmic
reticulum, or nuclear envelope (reviewed in Reed, J. C.,
Nature, 387:773-776 (1997); Adams & Cory, Science,
10 281:1322-1326 (1998); Gross et al., Genes Dev., 13:1899-
1911 (1999)). However, some pro-apoptotic Bcl-2 family
proteins, such as Bax, Bid, and Bim, are found in the
cytosol and must be induced to translocate to
mitochondria and other organelles where the Bcl-2-family
15 proteins to which they dimerize reside (Wolter et al., J.
Cell Biol., 139:1281-1292 (1997); Puthalakath et al.,
Mol. Cell, 3:287-96 (1999); Li et al., Cell, 94:491-501
(1998); Luo et al., Cell, 94:481-490 (1998)).

The intracellular locations of the Bcl-G_L and
20 Bcl-G_s protein was examined by confocal microscopy
analysis of cells expressing GFP-tagged proteins.
GFP-expressing cells were imaged by confocal microscopy
using a Bio-Rad MRC 1024 instrument (Xu & Reed, supra.
(1998); Zhang et al., supra. (2000); Zha et al., Mol.
25 Cell. Biol., 16:6494-6508 (1996)). GFP-Bcl-G_L protein was
located diffusely throughout cells, similar to GFP
control protein (Figure 11A, B). In contrast, Bcl-G_s was
found in a punctate cytosolic pattern (Figure 11C),
suggestive of organelle association. Surprisingly,
30 deletion of the BH3 domain from Bcl-G_s did not disrupt the
punctate distribution (Figure 5D), indicating that other
regions of the Bcl-G_s protein are sufficient for

subcellular targeting. Subcellular fractionation experiments confirmed these observations, demonstrating association of Bcl-G_s and Bcl-G_s (DB43) predominantly with organelle-containing heavy-membrane fractions, with scant
5 amounts in the soluble cytosolic compartment.

EXAMPLE VIII

Loss of Heterozygosity (LOH) is Associated with Bcl-G in Ovarian Tissue

This example describes loss of heterozygosity
10 (LOH) associated with Bcl-G in ovarian cancer tissue.

Ovarian cancer tissue samples were tested for SSCP for possible mutations in Bcl-G. No mutation was found in exon 1. However, about one third of the ovarian samples showed a possible LOH of Bcl-G. The LOH was
15 observed as a change in band intensity using SSCP. The results were confirmed independently using PCR. The LOH samples are sequenced to determine specific mutations.

These results indicate that LOH is associated with Bcl-G in ovarian tissue and can be useful as a
20 marker for ovarian cancer.

Example IX

Cloning of Mouse Bcl-G

This example describes cloning of mouse Bcl-G.

The mouse Bcl-G was identified by searching
25 GenBank. An EST clone (AA536718) was found to contain mouse Bcl-G. The EST was purchased from the American Type Culture Collection (ATCC; Manassas VA) and sequenced

to determine the complete sequence of mouse Bcl-G.

The nucleotide sequence of mouse Bcl-G cDNA is referenced as SEQ ID NO:41. The amino acid sequence of Bcl-G is referenced as SEQ ID NO:42.

5 PCR was used to isolate mouse Bcl-G from the
purchased EST clone and clone it into EGFP-C1 vector. he
primers used were MXSTA,
5'-GGGCTCGAGATGTGCAGCACCAGTGTGTATG-3' (SEQ ID NO:43);
NHREV, 5'-CCAAGCTTTAAGTCTACTTCTTCATGTGATATCCC-3' (SEQ ID
10 NO:44).

In preliminary experiments, mouse Bcl-G was overexpressed in Cos-7 and 293T cells. In these preliminary experiments, apoptosis was not observed.

15 Throughout this application various
publications have been referenced. The disclosures of
these publications in their entireties are hereby
incorporated by reference in this application in order to
more fully describe the state of the art to which this
invention pertains.

20 Although the invention has been described with
reference to the examples provided above, it should be
understood that various modifications can be made without
departing from the spirit of the invention. Accordingly,
the invention is limited only by the claims.

What is claimed is:

1. An isolated nucleic acid encoding a Bcl-G polypeptide, or a functional fragment thereof.

2. An isolated nucleic acid encoding a Bcl-G polypeptide, or a functional fragment thereof, comprising
5 a nucleic acid selected from:

(a) nucleic acid encoding the amino acid sequence set forth in SEQ ID NOS:2, 4 or 42, or

(b) nucleic acid that hybridizes to the
10 nucleic acid of (a) under moderately stringent conditions, wherein said nucleic acid contiguously encodes biologically active Bcl-G, or

(c) nucleic acid degenerate with respect to either (a) or (b) above, wherein said nucleic acid
15 encodes biologically active Bcl-G.

3. The nucleic acid of claim 2, wherein said nucleic acid hybridizes under high stringency conditions to the Bcl-G coding portion of any of SEQ ID NOS:1, 3 or 41.

20 4. The nucleic acid of claim 2, wherein the nucleotide sequence of said nucleic acid is substantially the same as set forth in any of SEQ ID NOS:1, 3 or 41.

5. The nucleic acid of claim 2, wherein the nucleotide sequence of said nucleic acid is the same as
25 that set forth in any of SEQ ID NOS:1, 3 or 41, or a modification thereof.

6. The nucleic acid of claim 2, wherein said nucleic acid is cDNA.

5 7. A vector containing the nucleic acid of claim 2.

8. Recombinant cells containing the nucleic acid of claim 2.

9. A Bcl-G oligonucleotide, comprising between 15 and 300 contiguous nucleotides of SEQ ID NOS:1, 3 or 10 41 or the anti-sense strand thereof.

10. An oligonucleotide according to claim 9, wherein said oligonucleotide is labeled with a detectable marker.

11. An antisense-nucleic acid capable of 15 specifically binding to mRNA encoded by said nucleic acid according to claim 2.

12. A kit for detecting the presence of a Bcl-G nucleic acid sequence comprising at least one oligonucleotide according to claim 10.

13. An isolated Bcl-G polypeptide, or a 20 functional fragment thereof, encoded by the nucleic acid of claim 2.

14. The Bcl-G polypeptide of claim 13, wherein said polypeptide comprises the substantially the same 25 amino acid sequence of Bcl-G_L (SEQ ID NO:2), Bcl-G_S (SEQ ID NO:4), or mouse Bcl-G (SEQ ID NO:42).

15. The Bcl-G polypeptide of claim 14, wherein said polypeptide comprises the amino acid sequence of Bcl-G_L (SEQ ID NO:2), Bcl-G_S (SEQ ID NO:4), or mouse Bcl-G (SEQ ID NO:42).

5 16. The Bcl-G polypeptide of claim 13, wherein said functional fragment comprises a BH3 or BH2 domain.

10 17. The Bcl-G polypeptide of claim 16, wherein said functional fragment comprises an amino acid sequence selected from the group consisting of SEQ ID NOS:5, 6, 9 and 18.

18. The Bcl-G polypeptide of claim 13, wherein said polypeptide is encoded by a nucleotide sequence comprising substantially the same nucleotide sequence as set forth in SEQ ID NOS:1, 3 or 41.

15 19. The Bcl-G polypeptide of claim 13, wherein said polypeptide is encoded by a nucleotide sequence comprising the sequence set forth in SEQ ID NOS:1 or 3.

20 20. A method for expression of a Bcl-G polypeptide, said method comprising culturing cells of claim 8 under conditions suitable for expression of said Bcl-G.

21. An isolated anti-Bcl-G antibody having specific reactivity with a Bcl-G according to claim 13.

25 22. Antibody according to claim 21, wherein said antibody is a monoclonal antibody.

23. A cell line producing the monoclonal antibody of claim 22.

24. An antibody according to claim 21, wherein said antibody is a polyclonal antibody.

25. A composition comprising an amount of the antisense-nucleic acid according to claim 11 effective to
5 inhibit expression of a human Bcl-G and an acceptable carrier capable of delivering Bcl-G to a cell.

26. A transgenic nonhuman mammal expressing exogenous nucleic acid according to claim 2, encoding a Bcl-G.

10 27. A transgenic nonhuman mammal according to claim 26, wherein said nucleic acid encoding said Bcl-G has been mutated, and wherein the Bcl-G so expressed is not native Bcl-G.

15 28. A mutant non-human mammal having a disrupted Bcl-G gene.

29. A transgenic nonhuman mammal according to claim 26, wherein the transgenic nonhuman mammal is a mouse.

20 30. A method for identifying nucleic acids encoding a mammalian Bcl-G, comprising contacting a sample containing nucleic acids with one or more oligonucleotides according to claim 9, wherein said contacting is effected under high stringency hybridization conditions, and identifying a nucleic acid
25 that hybridizes to said oligonucleotide.

31. A method of detecting a Bcl-G nucleic acid molecule in a sample, comprising contacting said sample with two or more Bcl-G oligonucleotides of claim 9, amplifying a nucleic acid molecule, and detecting said amplification.

32. The method of claim 31, wherein said amplification is performed using polymerase chain reaction.

33. A method for detecting the presence of a human Bcl-G in a sample, comprising contacting a sample with an antibody according to claim 21, and detecting the presence of specific binding of said antibody to said sample, thereby detecting the presence of a human Bcl-G in said sample.

34. Single stranded nucleic acid primers for amplification of a Bcl-G nucleic acid, wherein said primers comprise a nucleic acid sequence derived from the nucleic acid sequences set forth as SEQ ID NOS:1 or 3.

35. A method for modulating the activity of an oncogenic polypeptide, comprising contacting said oncogenic polypeptide with a substantially pure Bcl-G, or an oncogenic protein-binding fragment thereof.

36. A method of identifying an effective agent that alters the association of a Bcl-G with a Bcl-G associated polypeptide (BAP), comprising the steps of:

(a) contacting said Bcl-G and said BAP polypeptide, under conditions that allow said Bcl-G and BAP polypeptide to associate, with a compound; and

5 (b) detecting the altered association of said Bcl-G and BAP polypeptide, thereby identifying a compound that is an effective agent for altering the association of said Bcl-G with BAP.

37. The method of claim 36, wherein said compound is a drug.

38. The method of claim 36, wherein said compound is a polypeptide.

10 39. A method for modulating an activity mediated by a Bcl-G polypeptide, comprising contacting said Bcl-G polypeptide with an effective, modulating amount of an agent identified by claim 36.

15 40. The method of claim 39, wherein said modulated activity is the binding of Bcl-G to a Bcl-2 family member.

41. A method of modulating the level apoptosis in a cell, comprising the steps of:

20 (a) introducing a nucleic acid molecule encoding a Bcl-G into the cell; and

(b) expressing said Bcl-G in said cell, wherein the expression of said Bcl-G modulates apoptosis in said cell.

42. A method of modulating the level of apoptosis in a cell, comprising introducing an antisense nucleotide sequence into the cell, wherein said antisense nucleotide sequence specifically hybridizes to a nucleic acid molecule encoding a Bcl-G, wherein said hybridization reduces or inhibits the expression of said Bcl-G in said cell.

43. A therapeutic composition comprising a pharmaceutically acceptable carrier and a compound selected from the group consisting of a Bcl-G polypeptide, a functional fragment of said Bcl-G, a Bcl-G modulating compound identified according to claim 36, and an anti-Bcl-G antibody.

44. A method of treating a pathology characterized by abnormal cell proliferation, comprising administering an effective amount of the composition according to claim 43.

45. A method of diagnosing a pathology characterized by an increased or decreased level of a Bcl-G in a subject, comprising the steps of:

- (a) obtaining a test sample from the subject;
- (b) contacting said sample with an agent that can bind said Bcl-G under suitable conditions, wherein said conditions allow specific binding of said agent to said Bcl-G; and
- (c) comparing the amount of said specific binding in said test sample with the amount of specific binding in a control sample, wherein an increased or decreased amount of said

specific binding in said test sample as compared to said control sample is diagnostic of a pathology.

46. The method of claim 45, wherein said agent
5 is selected from the group consisting of an anti-Bcl-G antibody, a Bcl-G-associated-protein (BAP), and a Bcl-G nucleic acid.

47. A method of modulating the level of
apoptosis in a cell, comprising contacting the cell with
10 a compound that effectively alters the association of Bcl-G with a Bcl-G-associated-protein in the cell, or that effectively alters the activity of a Bcl-G in the cell.

48. A chimeric protein comprising a domain
15 selected from the group consisting of BH3 (SEQ ID NOS:5 or 9) and BH2 (SEQ ID NOS:6 or 18).

49. A method of modulating interactions
between Bcl-G and Bcl-2, comprising contacting a Bcl-G
polypeptide with the agent of claim 36.

20 50. A method of diagnosing cancer or monitoring cancer therapy comprising contacting a test sample from a patient with the antibody of claim 21.

51. A method of assessing prognosis of
patients with cancer comprising contacting a test sample
25 from a patient with the antibody of claim 21.

52. A method of diagnosing cancer or monitoring cancer therapy comprising contacting a test sample from a patient with the oligonucleotide of claim 9.

- 5 53. A method of assessing prognosis of patients with cancer comprising contacting a test sample from a patient with the oligonucleotide of claim 9.

```

      10      20      30      40
      |-----|
aatgacatgacagccattccgtggccagggacaccactgc 40
ccaagctggagaccacgaggattcagggactgaagccagc 80
atgggaattcctgggtttgagatcagagtcctgagtacctc 120
gtgggaacttgggcactcatccgcaggaggtctagacccc 160
caggaattccttgagtctaaggcacaggcccaacatgtg 200

      210      220      230      240
      |-----|
tagcaccagtgggtgtgacctggaagaaatccccctagat 240
gatgatgacctaaacaccatagaattcaaaatcctcgcc 280
actacaccagacatcatgtcttcaagagcaccctgtct 320
cttctcaccaaagctgtgagaacaagaagtttgtccag 360
aggggcctggggaattgttcagcaaatgagtcatggacag 400

      410      420      430      440
      |-----|
aggtgtcatggccttgcagaaattcccaatccagtggagaa 440
ggccataaaccttggcaagaaaaagtcttcttggaaagca 480
ttctttggagtagtggagaagggaagattcgcagagcacgc 520
ctgccaaaggtctctgtctcagggtcaaaggacgttggata 560
ccaagattcgcacagccagcagtggtccagggtgtcttct 600

      610      620      630      640
      |-----|
aacgtggagcagtgcttggagcatgaagctgtggaccca 640
aagtcatttccattgccaaaccgagtagctgaaattgttta 680
ctcctggccaccaccacaagcgacccaggcaggaggcttc 720
aagtccaaagagatttttgaactgagggtctctccttcc 760
agtcctcaaggccacgtgcctgtogcttcaagttctaagaa 800

      810      820      830      840
      |-----|
agatgaagaagaacaaatactagccaaaattgttgagctg 840
ctgaaatattcaggagatcagttggaagaaagctgaaga 880
aagataaggctttgatgggccacttccaggatgggctgtc 920
ctactctgttttcaagaccatcacagaccaggtcctaag 960
gggtgtggacccaggggagaatcagaggtcaaagctcagg 1000

      1010      1020      1030      1040
      |-----|
gctttaaggctgcccttgaatagacgtcacggccaagct 1040
cacagctattgacaaccaccgatgaacagggtcctgggc 1080
tttggcaccaagtacctgaaagagaacttctcgccatgga 1120
tccagcagcacgggtggatgggaaaaaatacttgggatatc 1160
acatgaagaagtagactga 1179

```

FIGURE 1

Bcl-G_L

1	- ATGTGTAGCAACAGTGGGTGTGACCTGGAAGAAATCCCCCTAGATGATGATGACCTAAAC	- 60
	- M C S T S G C D L E E I P L D D D D L N	
61	- ACCATAGAATTCAAAATCCTGGCTACTACACCAGACATCATGTCTTCAAGAGCACCCCT	- 120
	- T I E F K I L A Y Y T R H H V F K S T P	
121	- GCTCTCTTCTCAACCAAGCTGCTGAGAACAAGAAGTTTGTGCCAGAGGGGCTGGGGAAT	- 180
	- A L F S P K L L R T R S L S Q R G L G N	
181	- TGTTCAGCAAAATGAGTCATGGACAGAGGTGTCTATGGGCTTGCAGAAATCCCAATCCAGT	- 240
	- C S A N E S W T E V S W P C R N S Q S S	
241	- GAGAAGGCCATAAACCTTGGCAAGAAAAGTCTTCTTGGAAAGCATCTCTTGGAGTAGTIG	- 300
	- E K A I N L G K K K S S W K A F F G V V	
301	- GAGAAGGAAGATTGGCAGAGCAAGGCTGCCAAGGTCTCTGCTCAGGGTCAAAGGAAGTTG	- 360
	- E K E D S Q S T P A K V S A Q G Q R T L	
361	- GAATACCAAGATTGGCAGAGCAGAGTGGTCCAGGTGTCTTCTTACGGTGGAGCAGTGC	- 420
	- E Y Q D S H S Q Q W S R C L S N V E Q C	
421	- TTGGAGCATGAAGCTGTGGACCCCAAAGTCATTTCATTTGCCAAGAGTAGCTGAATTT	- 480
	- L E H E A V D P K V I S I A N R V A E I	
481	- GTTACTCTCTGGCCACCAACCAAGGACCCAGGCAGGAGCTTCAAGTCCAAAGAGATT	- 540
	- V Y S W P P P Q A T Q A G G F K S K E I	
541	- TTGTAACTGAGGGTCTCTCTCTTCCAGCTCCAAGGCCAGTCCCTGTAGCTTCAAGTTCT	- 600
	- F V T E G L S F Q L Q G H V P V A S S S	
601	- AAGAAAGATGAAGAAGAACAAATACCTAGCCAAATTTGTGAGCTGCTGAAATATTTCAGGA	- 660
	- K K D E E E Q I L A K I V E L L K Y S G	
661	- GATCAGTTGGAAAGAAAGCTGAAGAAAGATAAGGCTTTGATGGGCCACTTCCAGGATGGG	- 720
	- D Q L E R K L K K D K A L M G H F Q D G	
721	- CTGTCTACTCTGTTTTCAAGACCATCACAGACAGGTCTTAATGGGTGTGGACCCAGG	- 780
	- L S Y S V F K T I T D Q V L M G V D P R	
781	- GGAGAATCAGAGGTCAAAGCTCAGGCTTTAAGGCTGGGCTTGTAAATAGAGTCAAGGCC	- 840
	- G E S E V K A Q G F K A A L V I D V T A	
841	- AAGCTCACAGCTATTGACAAACCCGATGAACAGGGTCTTGGGCTTTGGCACCAAGTAC	- 900
	- K L T A I D N H P M N R V L G F G T K Y	
901	- CTGAAAGAGAACTTCTGGCATGGATCCAGCAGCAAGGTGGATGGGAAAAATACTTGGG	- 960
	- L K E N F S P W I Q Q H G G W E K I L G	
961	- ATATCATGAAGAAGTAGACTGA	- 984
	- I S H E E V D *	

FIGURE 2

```

      10      20      30      40
aatgacatgacagccattccgtggccagggacaccactgc 40
ccaagctggagaccacgaggattcagggactgaagccagc 80
atgggaattcctggtttgagatcagagtcctgagtacctc 120
gtgggaacttgggcactcatccgcaggaggcttagacccc 160
cagagaattccttgagcttaaggcacaggcccaacatgtg 200

      210      220      230      240
tagcaccagtggggtgtgacctggaagaaatccccctagat 240
gatgatgacctaaacaccatagaattcaaaatcctcgct 280
actacaccagacatcatgtcttcaagagcaccctgtct 320
cttctcaccaaagctgctgagaacaagaagttgtcccag 360
aggggcctggggaattgttcagcaaatgagtcattggacag 400

      410      420      430      440
aggtgtcatggccttgacagaaattcccaatccagtgagaa 440
ggccataaaccttggcaagaaaaagctcttcttggaagca 480
ttctttggagtagtggagaaggagattcgacagacacgc 520
ctgccaaaggtctctgctcaggggtcaaaggacgttggaata 560
ccaagattcgacagccagcagtggtccaggtgtctttct 600

      610      620      630      640
aacgtggagcagtgcttggagcatgaagctgtggacccca 640
aagtcatttccattgccaaccgagtagctgaaattgttta 680
ctcctggccaccaccacaagcgacccaggcaggaggcttc 720
aagtccaaagagatttttgtaactgaggggtctctccttc 760
agctccaaggccacgtgcctgtagcttcaagttctaagaa 800

      810      820      830      840
agatgaagaagaacaaatactagccaaaattggtgagctg 840
ctgaaatattcaggagatcagttggaaagaaaggacactg 880
ccttcatccccattcccttgggtgacaccagcatccaggg 920
ttttccacaggatgggttgatggcctgcatttga 954

```

FIGURE 3

Bcl-Gs

```

1 - ATGTGTAGCACAGTGGGTGTGACCTGGAAGAAATCCCCCTAGATGATGACCTAAAC - 60
  - M C S T S G C D L E E I P L D D D L N
61 - ACCATAGNATTCAAATCTCGCTACTACACACATCATCTCTTCAGAGACCCCT - 120
  - T I E F K I L A Y Y T R H V F K S T P
121 - GCTCTCTCTCACCAAGCTGCTCAGAACAGAGTTTGTCCAGAGGGGCTGGGAAT - 180
  - A L F S P K L L R T R S L S Q R G L G N
181 - TGTTCAGCAATGATGATGACAGAGGTGTCATGGCTTGCAGAAATCCCAATCCAGT - 240
  - C S A N E S W T E V S W P C R N S Q S S
241 - GAGAAGGCCATAAACCTTGCAAGAAAGTCTCTTGAAGACATCTTTGGAGTAGTG - 300
  - E K A I N L G K K K S S W K A F F G V V
301 - GAGAAGGAAGATTCCGAGAGCACCCCTCCCAAGGTCTCTGCTCAGGTCAAAGACGTTG - 360
  - E K E D S Q S T P A K V S A Q G Q R T L
361 - GAATACCAAGATTCCACAGCCAGCGAGTGTCAGGTGTCTTTCTACGTGAGCAGTGC - 420
  - E Y Q D S H S Q Q W S R C L S N V E Q C
421 - TTGGAGCATGAAGCTGTGACCCCAAGTCAATTCCTCCACCGAGTAGCTGAAT - 480
  - L E H E A V D P K V I S I A N R V A E I
481 - GTTTACTCTGCCACCACACAGCCAGCCAGCGAGGCTTCAAGTCCAAAGAGATT - 540
  - V Y S W P P P Q A T Q A G G F K S K E I
541 - TTGTAACTGAGGGTCTCTCTTCAGCTCCAGGCCAGCTGCTGTAGCTTCAAGTTCT - 600
  - F V T E G L S F Q L Q G H V P V A S S S
601 - AAGAAGATGAAGAAGAACAAATACTAGCCAAATTTGTGAGCTGTGAAATATTCAGGA - 660
  - K K D E E E Q I L A K I V E L L K Y S G
661 - GATCAGTTGGAAGAAAGACACTGCCTTCATCCCATTCCTTGTGACACCAATC - 720
  - D L E R K D T A F I P I P L V D T S I
721 - CAGGGTTTCCACAGGATGTTGATGCGCTGCATTGA - 759
  - Q G F P Q D G L M A C I .

```

FIGURE 4




FIGURE 5




FIGURE 6





FIGURE 7

**FIGURE 8**




FIGURE 9




FIGURE 10




FIGURE 11

SEQUENCE LISTING

<110> Reed, John C.
Godzik, Adam
The Burnham Institute

<120> BCL-G Polypeptides, Encoding Nucleic Acids and Methods
of Use

<130> FP-LJ 4352

<140>

<141>

<150> US 09/461,641

<151> 1999-12-14

<160> 44

<170> PatentIn Ver. 2.0

<210> 1

<211> 1179

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (196)..(1179)

<400> 1

```

aatgacatga cagccattcc gtggccaggg acaccactgc ccaagctgga gaccacgagg 60
attcagggaac tgaagccagc atgggaattc ctggtttgag atcagagtcc tgagtacctc 120
gtgggaactt gggcactcat ccgcaggagg tctagacccc cagagaattc cttgagtcta 180
aggcacaggc ccaac atg tgt agc acc agt ggg tgt gac ctg gaa gaa atc 231
      Met Cys Ser Thr Ser Gly Cys Asp Leu Glu Glu Ile
          1             5             10

ccc cta gat gat gat gac cta aac acc ata gaa ttc aaa atc ctc gcc 279
Pro Leu Asp Asp Asp Asp Leu Asn Thr Ile Glu Phe Lys Ile Leu Ala
      15             20             25

tac tac acc aga cat cat gtc ttc aag agc acc cct gct ctc ttc tca 327
Tyr Tyr Thr Arg His His Val Phe Lys Ser Thr Pro Ala Leu Phe Ser
      30             35             40

cca aag ctg ctg aga aca aga agt ttg tcc cag agg ggc ctg ggg aat 375
Pro Lys Leu Leu Arg Thr Arg Ser Leu Ser Gln Arg Gly Leu Gly Asn
      45             50             55             60

tgt tca gca aat gag tca tgg aca gag gtg tca tgg cct tgc aga aat 423
Cys Ser Ala Asn Glu Ser Trp Thr Glu Val Ser Trp Pro Cys Arg Asn
      65             70             75

```

tcc caa tcc agt gag aag gcc ata aac ctt ggc aag aaa aag tct tct	471
Ser Gln Ser Ser Glu Lys Ala Ile Asn Leu Gly Lys Lys Lys Ser Ser	
80 85 90	
tgg aaa gca ttc ttt gga gta gtg gag aag gaa gat tcg cag agc acg	519
Trp Lys Ala Phe Phe Gly Val Val Glu Lys Glu Asp Ser Gln Ser Thr	
95 100 105	
cct gcc aag gtc tct gct cag ggt caa agg acg ttg gaa tac caa gat	567
Pro Ala Lys Val Ser Ala Gln Gly Gln Arg Thr Leu Glu Tyr Gln Asp	
110 115 120	
tcg cac agc cag cag tgg tcc agg tgt ctt tct aac gtg gag cag tgc	615
Ser His Ser Gln Gln Trp Ser Arg Cys Leu Ser Asn Val Glu Gln Cys	
125 130 135 140	
ttg gag cat gaa gct gtg gac ccc aaa gtc att tcc att gcc aac cga	663
Leu Glu His Glu Ala Val Asp Pro Lys Val Ile Ser Ile Ala Asn Arg	
145 150 155	
gta gct gaa att gtt tac tcc tgg cca cca cca caa gcg acc cag gca	711
Val Ala Glu Ile Val Tyr Ser Trp Pro Pro Pro Gln Ala Thr Gln Ala	
160 165 170	
gga ggc ttc aag tcc aaa gag att ttt gta act gag ggt ctc tcc ttc	759
Gly Gly Phe Lys Ser Lys Glu Ile Phe Val Thr Glu Gly Leu Ser Phe	
175 180 185	
cag ctc caa ggc cac gtg cct gta gct tca agt tct aag aaa gat gaa	807
Gln Leu Gln Gly His Val Pro Val Ala Ser Ser Ser Lys Lys Asp Glu	
190 195 200	
gaa gaa caa ata cta gcc aaa att gtt gag ctg ctg aaa tat tca gga	855
Glu Glu Gln Ile Leu Ala Lys Ile Val Glu Leu Leu Lys Tyr Ser Gly	
205 210 215 220	
gat cag ttg gaa aga aag ctg aag aaa gat aag gct ttg atg ggc cac	903
Asp Gln Leu Glu Arg Lys Leu Lys Lys Asp Lys Ala Leu Met Gly His	
225 230 235	
ttc cag gat ggg ctg tcc tac tct gtt ttc aag acc atc aca gac cag	951
Phe Gln Asp Gly Leu Ser Tyr Ser Val Phe Lys Thr Ile Thr Asp Gln	
240 245 250	
gtc cta atg ggt gtg gac ccc agg gga gaa tca gag gtc aaa gct cag	999
Val Leu Met Gly Val Asp Pro Arg Gly Glu Ser Glu Val Lys Ala Gln	
255 260 265	
ggc ttt aag gct gcc ctt gta ata gac gtc acg gcc aag ctc aca gct	1047
Gly Phe Lys Ala Ala Leu Val Ile Asp Val Thr Ala Lys Leu Thr Ala	
270 275 280	
att gac aac cac ccg atg aac agg gtc ctg ggc ttt ggc acc aag tac	1095
Ile Asp Asn His Pro Met Asn Arg Val Leu Gly Phe Gly Thr Lys Tyr	
285 290 295 300	
ctg aaa gag aac ttc tcg cca tgg atc cag cag cac ggt gga tgg gaa	1143

Leu Lys Glu Asn Phe Ser Pro Trp Ile Gln Gln His Gly Gly Trp Glu
 305 310 315

aaa ata ctt ggg ata tca cat gaa gaa gta gac tga 1179
 Lys Ile Leu Gly Ile Ser His Glu Glu Val Asp
 320 325

<210> 2
 <211> 327
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Cys Ser Thr Ser Gly Cys Asp Leu Glu Glu Ile Pro Leu Asp Asp
 1 5 10 15

Asp Asp Leu Asn Thr Ile Glu Phe Lys Ile Leu Ala Tyr Tyr Thr Arg
 20 25 30

His His Val Phe Lys Ser Thr Pro Ala Leu Phe Ser Pro Lys Leu Leu
 35 40 45

Arg Thr Arg Ser Leu Ser Gln Arg Gly Leu Gly Asn Cys Ser Ala Asn
 50 55 60

Glu Ser Trp Thr Glu Val Ser Trp Pro Cys Arg Asn Ser Gln Ser Ser
 65 70 75 80

Glu Lys Ala Ile Asn Leu Gly Lys Lys Lys Ser Ser Trp Lys Ala Phe
 85 90 95

Phe Gly Val Val Glu Lys Glu Asp Ser Gln Ser Thr Pro Ala Lys Val
 100 105 110

Ser Ala Gln Gly Gln Arg Thr Leu Glu Tyr Gln Asp Ser His Ser Gln
 115 120 125

Gln Trp Ser Arg Cys Leu Ser Asn Val Glu Gln Cys Leu Glu His Glu
 130 135 140

Ala Val Asp Pro Lys Val Ile Ser Ile Ala Asn Arg Val Ala Glu Ile
 145 150 155 160

Val Tyr Ser Trp Pro Pro Pro Gln Ala Thr Gln Ala Gly Gly Phe Lys
 165 170 175

Ser Lys Glu Ile Phe Val Thr Glu Gly Leu Ser Phe Gln Leu Gln Gly
 180 185 190

His Val Pro Val Ala Ser Ser Ser Lys Lys Asp Glu Glu Glu Gln Ile
 195 200 205

Leu Ala Lys Ile Val Glu Leu Leu Lys Tyr Ser Gly Asp Gln Leu Glu
 210 215 220

Arg Lys Leu Lys Lys Asp Lys Ala Leu Met Gly His Phe Gln Asp Gly

```

225                230                235                240
Leu Ser Tyr Ser Val Phe Lys Thr Ile Thr Asp Gln Val Leu Met Gly
                245                250                255
Val Asp Pro Arg Gly Glu Ser Glu Val Lys Ala Gln Gly Phe Lys Ala
                260                265                270
Ala Leu Val Ile Asp Val Thr Ala Lys Leu Thr Ala Ile Asp Asn His
                275                280                285
Pro Met Asn Arg Val Leu Gly Phe Gly Thr Lys Tyr Leu Lys Glu Asn
                290                295                300
Phe Ser Pro Trp Ile Gln Gln His Gly Gly Trp Glu Lys Ile Leu Gly
305                310                315                320
Ile Ser His Glu Glu Val Asp
                325

```

```

<210> 3
<211> 954
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> CDS
<222> (196)..(954)

```

```

<400> 3
aatgacatga cagccattcc gtggccaggg acaccactgc ccaagctgga gaccacgagg 60
attcagggac tgaagccagc atgggaattc ctggtttgag atcagagtcc tgagtacctc 120
gtgggaactt gggcactcat ccgcaggagg tctagacccc cagagaattc cttgagtcta 180
aggcacaggc ccaac atg tgt agc acc agt ggg tgt gac ctg gaa gaa atc 231
      Met Cys Ser Thr Ser Gly Cys Asp Leu Glu Ile
              1              5              10
ccc cta gat gat gat gac cta aac acc ata gaa ttc aaa atc ctc gcc 279
Pro Leu Asp Asp Asp Asp Leu Asn Thr Ile Glu Phe Lys Ile Leu Ala
      15              20              25
tac tac acc aga cat cat gtc ttc aag agc acc cct gct ctc ttc tca 327
Tyr Tyr Thr Arg His His Val Phe Lys Ser Thr Pro Ala Leu Phe Ser
      30              35              40
cca aag ctg ctg aga aca aga agt ttg tcc cag agg ggc ctg ggg aat 375
Pro Lys Leu Leu Arg Thr Arg Ser Leu Ser Gln Arg Gly Leu Gly Asn
      45              50              55              60
tgt tca gca aat gag tca tgg aca gag gtg tca tgg cct tgc aga aat 423
Cys Ser Ala Asn Glu Ser Trp Thr Glu Val Ser Trp Pro Cys Arg Asn
      65              70              75

```

tcc caa tcc agt gag aag gcc ata aac ctt ggc aag aaa aag tct tct	471
Ser Gln Ser Ser Glu Lys Ala Ile Asn Leu Gly Lys Lys Lys Ser Ser	
80 85 90	
tggtg aaa gca ttc ttt gga gta gtg gag aag gaa gat tcg cag agc acg	519
Trp Lys Ala Phe Phe Gly Val Val Glu Lys Glu Asp Ser Gln Ser Thr	
95 100 105	
cct gcc aag gtc tct gct cag ggt caa agg acg ttg gaa tac caa gat	567
Pro Ala Lys Val Ser Ala Gln Gly Gln Arg Thr Leu Glu Tyr Gln Asp	
110 115 120	
tcg cac agc cag cag tgg tcc agg tgt ctt tct aac gtg gag cag tgc	615
Ser His Ser Gln Gln Trp Ser Arg Cys Leu Ser Asn Val Glu Gln Cys	
125 130 135 140	
ttg gag cat gaa gct gtg gac ccc aaa gtc att tcc att gcc aac cga	663
Leu Glu His Glu Ala Val Asp Pro Lys Val Ile Ser Ile Ala Asn Arg	
145 150 155	
gta gct gaa att gtt tac tcc tgg cca cca cca caa gcg acc cag gca	711
Val Ala Glu Ile Val Tyr Ser Trp Pro Pro Pro Gln Ala Thr Gln Ala	
160 165 170	
gga ggc ttc aag tcc aaa gag att ttt gta act gag ggt ctc tcc ttc	759
Gly Gly Phe Lys Ser Lys Glu Ile Phe Val Thr Glu Gly Leu Ser Phe	
175 180 185	
cag ctc caa ggc cac gtg cct gta gct tca agt tct aag aaa gat gaa	807
Gln Leu Gln Gly His Val Pro Val Ala Ser Ser Ser Lys Lys Asp Glu	
190 195 200	
gaa gaa caa ata cta gcc aaa att gtt gag ctg ctg aaa tat tca gga	855
Glu Glu Gln Ile Leu Ala Lys Ile Val Glu Leu Leu Lys Tyr Ser Gly	
205 210 215 220	
gat cag ttg gaa aga aag gac act gcc ttc atc ccc att ccc ttg gtt	903
Asp Gln Leu Glu Arg Lys Asp Thr Ala Phe Ile Pro Ile Pro Leu Val	
225 230 235	
gac acc agc atc cag ggt ttt cca cag gat ggt ttg atg gcc tgc att	951
Asp Thr Ser Ile Gln Gly Phe Pro Gln Asp Gly Leu Met Ala Cys Ile	
240 245 250	
tga	954

<210> 4
 <211> 252
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Cys Ser Thr Ser Gly Cys Asp Leu Glu Glu Ile Pro Leu Asp Asp
 1 5 10 15

Asp Asp Leu Asn Thr Ile Glu Phe Lys Ile Leu Ala Tyr Tyr Thr Arg
 20 25 30
 His His Val Phe Lys Ser Thr Pro Ala Leu Phe Ser Pro Lys Leu Leu
 35 40 45
 Arg Thr Arg Ser Leu Ser Gln Arg Gly Leu Gly Asn Cys Ser Ala Asn
 50 55 60
 Glu Ser Trp Thr Glu Val Ser Trp Pro Cys Arg Asn Ser Gln Ser Ser
 65 70 75 80
 Glu Lys Ala Ile Asn Leu Gly Lys Lys Lys Ser Ser Trp Lys Ala Phe
 85 90 95
 Phe Gly Val Val Glu Lys Glu Asp Ser Gln Ser Thr Pro Ala Lys Val
 100 105 110
 Ser Ala Gln Gly Gln Arg Thr Leu Glu Tyr Gln Asp Ser His Ser Gln
 115 120 125
 Gln Trp Ser Arg Cys Leu Ser Asn Val Glu Gln Cys Leu Glu His Glu
 130 135 140
 Ala Val Asp Pro Lys Val Ile Ser Ile Ala Asn Arg Val Ala Glu Ile
 145 150 155 160
 Val Tyr Ser Trp Pro Pro Pro Gln Ala Thr Gln Ala Gly Gly Phe Lys
 165 170 175
 Ser Lys Glu Ile Phe Val Thr Glu Gly Leu Ser Phe Gln Leu Gln Gly
 180 185 190
 His Val Pro Val Ala Ser Ser Ser Lys Lys Asp Glu Glu Glu Gln Ile
 195 200 205
 Leu Ala Lys Ile Val Glu Leu Leu Lys Tyr Ser Gly Asp Gln Leu Glu
 210 215 220
 Arg Lys Asp Thr Ala Phe Ile Pro Ile Pro Leu Val Asp Thr Ser Ile
 225 230 235 240
 Gln Gly Phe Pro Gln Asp Gly Leu Met Ala Cys Ile
 245 250

<210> 5

<211> 9

<212> PRT

<213> Homo sapiens

<400> 5

Leu Lys Tyr Ser Gly Asp Gln Leu Glu
 1 5

<210> 6

<211> 10
 <212> PRT
 <213> Homo sapiens

<400> 6
 Pro Trp Ile Gln Gln His Gly Gly Trp Glu
 1 5 10

<210> 7
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 7
 gtacttggtg ccaaagccca gg 22

<210> 8
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 8
 gacatgatgt ctggtgtagt aggcgagg 28

<210> 9
 <211> 15
 <212> PRT
 <213> Homo sapiens

<400> 9
 Ile Val Glu Leu Leu Lys Tyr Ser Gly Asp Gln Leu Glu Arg Lys
 1 5 10 15

<210> 10
 <211> 15
 <212> PRT
 <213> Homo sapiens

<400> 10
 Leu Ser Glu Cys Leu Lys Arg Ile Gly Asp Glu Leu Asp Ser Asn
 1 5 10 15

<210> 11
 <211> 15
 <212> PRT
 <213> Homo sapiens

<400> 11

Val	Gly	Arg	Gln	Leu	Ala	Ile	Ile	Gly	Asp	Asp	Ile	Asn	Arg	Arg
1				5					10					15

<210> 12

<211> 15

<212> PRT

<213> Homo sapiens

<400> 12

Ile	Ala	Arg	His	Leu	Ala	Gln	Val	Gly	Asp	Ser	Met	Asp	Arg	Ser
1				5					10					15

<210> 13

<211> 15

<212> PRT

<213> Homo sapiens

<400> 13

Leu	Ala	Leu	Arg	Leu	Ala	Cys	Ile	Gly	Asp	Glu	Met	Asp	Val	Ser
1				5					10					15

<210> 14

<211> 15

<212> PRT

<213> Homo sapiens

<400> 14

Ala	Leu	Glu	Thr	Leu	Arg	Arg	Val	Gly	Asp	Gly	Val	Gln	Arg	Asn
1				5					10					15

<210> 15

<211> 15

<212> PRT

<213> Homo sapiens

<400> 15

Val	Lys	Gln	Ala	Leu	Arg	Glu	Ala	Gly	Asp	Glu	Phe	Glu	Leu	Arg
1				5					10					15

<210> 16

<211> 15

<212> PRT

<213> Homo sapiens

<400> 16

Leu	His	Gln	Ala	Met	Arg	Ala	Ala	Gly	Asp	Glu	Phe	Glu	Thr	Arg
1				5					10					15

<210> 17

<211> 15
<212> PRT
<213> Homo sapiens

<400> 17
Val His Leu Thr Leu Arg Gln Ala Gly Asp Asp Phe Ser Arg Arg
1 5 10 15

<210> 18
<211> 8
<212> PRT
<213> Homo sapiens

<400> 18
Trp Ile Gln Gln His Gly Gly Trp
1 5

<210> 19
<211> 8
<212> PRT
<213> Homo sapiens

<400> 19
Trp Ile Gln Asp Gln Gly Gly Trp
1 5

<210> 20
<211> 8
<212> PRT
<213> Homo sapiens

<400> 20
Trp Ile Ala Gln Arg Gly Gly Trp
1 5

<210> 21
<211> 8
<212> PRT
<213> Homo sapiens

<400> 21
Trp Leu Arg Arg Arg Gly Gly Trp
1 5

<210> 22
<211> 8
<212> PRT
<213> Homo sapiens

<400> 22
Trp Ile Arg Gln Asn Gly Gly Trp
1 5

<210> 23
<211> 8
<212> PRT
<213> Homo sapiens

<400> 23
Trp Leu Val Lys Gln Arg Gly Trp
1 5

<210> 24
<211> 8
<212> PRT
<213> Homo sapiens

<400> 24
Trp Ile Gln Glu Asn Gly Gly Trp
1 5

<210> 25
<211> 8
<212> PRT
<213> Homo sapiens

<400> 25
Trp Ile His Ser Ser Gly Gly Trp
1 5

<210> 26
<211> 8
<212> PRT
<213> Homo sapiens

<400> 26
Trp Ile Gln Asp Asn Gly Gly Trp
1 5

<210> 27
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 27
ggctcgagcg atgtgtagca ccagtgggtg tgacc

35

<210> 28
<211> 35

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 28
ccaagcttta agtctacttc ttcattgtgat atccc 35

<210> 29
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 29
ccaagcttta aaatgcaggc catcaaacc 29

<210> 30
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 30
ggctcgagcg atgtgtagca ccagtgggtg tgacc 35

<210> 31
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 31
ccggatccgg ctagtatttg ttcttcttca tcttt 35

<210> 32
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic

primer

<400> 32
ccggatccga cactgccttc atccccattc cc 32

<210> 33
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic
primer

<400> 33
ccaagcttta aaatgcaggc catcaaacc 29

<210> 34
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 34
gccaaaattg ttgagctgga gaaatattca ggagatcagt tgg 43

<210> 35
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 35
ccaactgac tcctgaatat ttctccagct caacaatttt ggc 43

<210> 36
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 36
ctgagggtct ctcttccag ctccaagg 28

<210> 37
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 37
ggccgtgacg tctattacaa gggcagcc

28

<210> 38
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 38
caagggaatg gggatgaagg cagtgtc

27

<210> 39
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 39
tgaaggtcgg agtcaacgga ttggt

26

<210> 40
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 40
catgtgggcc atgaggtcca ccac

24

<210> 41
<211> 987
<212> DNA
<213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(987)

<220>
 <221> unsure
 <222> (319)
 <223> unknown nucleotide

<400> 41
 atg tgc agc acc agt gtg tat gac ctg gaa gac att ccc ctg gag gat 48
 Met Cys Ser Thr Ser Val Tyr Asp Leu Glu Asp Ile Pro Leu Glu Asp
 1 5 10 15
 gat gat cca aac agc ata gag ttc aaa atc ctg gcc ttc tac gcc aga 96
 Asp Asp Pro Asn Ser Ile Glu Phe Lys Ile Leu Ala Phe Tyr Ala Arg
 20 25 30
 cac cat gtc ttc aag aac acc ccg gct gtc ttc tcg ccc aag ctc tcc 144
 His His Val Phe Lys Asn Thr Pro Ala Val Phe Ser Pro Lys Leu Ser
 35 40 45
 aga aca agg agt ctg tcc cag aaa gcc ctg ggg act tgg tca act gat 192
 Arg Thr Arg Ser Leu Ser Gln Lys Ala Leu Gly Thr Trp Ser Thr Asp
 50 55 60
 tcc tgg aca cag gta tca ttg cct tgc aga ggt tcc ccc tcc agc gaa 240
 Ser Trp Thr Gln Val Ser Leu Pro Cys Arg Gly Ser Pro Ser Ser Glu
 65 70 75 80
 aag aac atc agc ttg ggc aag aag aag tct tct tgg aga aca ctc ttc 288
 Lys Asn Ile Ser Leu Gly Lys Lys Lys Ser Ser Trp Arg Thr Leu Phe
 85 90 95
 agg gtg gcc gag aag gag gaa ggc ctg ccg ngc tcc cca aag gag atc 336
 Arg Val Ala Glu Lys Glu Glu Gly Leu Pro Xaa Ser Pro Lys Glu Ile
 100 105 110
 cga gct cag ggt cct cag ggc ccc ttc ccg gta gag cgg cag agt ggc 384
 Arg Ala Gln Gly Pro Gln Gly Pro Phe Pro Val Glu Arg Gln Ser Gly
 115 120 125
 ttc cac aac cag cac tgg ccc agg tct ctg agc agt gtg gag cag ccc 432
 Phe His Asn Gln His Trp Pro Arg Ser Leu Ser Ser Val Glu Gln Pro
 130 135 140
 tgg aga gtg aag ttg tgg att cca aag tgg ctt gta ttg cca aca gag 480
 Trp Arg Val Lys Leu Trp Ile Pro Lys Trp Leu Val Leu Pro Thr Glu
 145 150 155 160
 tgg ctg aaa ttg ttt act cct ggc cac cac cag atg tca tcc aca gcc 528
 Trp Leu Lys Leu Phe Thr Pro Gly His His Gln Met Ser Ser Thr Ala
 165 170 175
 agg gag gaa gcc agc tca aag aga ggg tct cgg aga ttt ttg tac ttc 576
 Arg Glu Glu Ala Ser Ser Lys Arg Gly Ser Arg Arg Phe Leu Tyr Phe

180	185	190	
agg ttt gaa gga cct tgg gac tct aag aat aaa gat ggt gaa gac caa			624
Arg Phe Glu Gly Pro Trp Asp Ser Lys Asn Lys Asp Gly Glu Asp Gln			
195	200	205	
ata ata agc aag att gtt gag ctg ctg aaa tcc tcg ggg gat cag ttg			672
Ile Ile Ser Lys Ile Val Glu Leu Leu Lys Ser Ser Gly Asp Gln Leu			
210	215	220	
gga aga gag ata aag aaa gac aag gct ttg atg agc agc ttc cag gac			720
Gly Arg Glu Ile Lys Lys Asp Lys Ala Leu Met Ser Ser Phe Gln Asp			
225	230	235	240
ggg ctg tcc tac tca acg ttc aag acc atc aca gac ctg ttc ctg agg			768
Gly Leu Ser Tyr Ser Thr Phe Lys Thr Ile Thr Asp Leu Phe Leu Arg			
245	250	255	
gac gtg gac acc aga gga gaa tca gag gtc aaa gct cgg ggc ttc aag			816
Asp Val Asp Thr Arg Gly Glu Ser Glu Val Lys Ala Arg Gly Phe Lys			
260	265	270	
gct gcc ctt gca ata gac gcc atc gcc aag ctc acg gca tcg gac aac			864
Ala Ala Leu Ala Ile Asp Ala Ile Ala Lys Leu Thr Ala Ser Asp Asn			
275	280	285	
cac cca atg aat aga atg ctg ggc ttc ggg acc aag tac cta aaa gag			912
His Pro Met Asn Arg Met Leu Gly Phe Gly Thr Lys Tyr Leu Lys Glu			
290	295	300	
tac ttc tcc ccc tgg gtt cag cag aat ggc gga tgg gaa aaa ata ctt			960
Tyr Phe Ser Pro Trp Val Gln Gln Asn Gly Gly Trp Glu Lys Ile Leu			
305	310	315	320
ggg atc tca cat gaa gaa gta gac tga			987
Gly Ile Ser His Glu Glu Val Asp			
325			

<210> 42
 <211> 328
 <212> PRT
 <213> Mus musculus

<220>
 <221> 107
 <222> unsure
 <223> Xaa = unknown amino acid

<400> 42
 Met Cys Ser Thr Ser Val Tyr Asp Leu Glu Asp Ile Pro Leu Glu Asp
 1 5 10 15
 Asp Asp Pro Asn Ser Ile Glu Phe Lys Ile Leu Ala Phe Tyr Ala Arg
 20 25 30
 His His Val Phe Lys Asn Thr Pro Ala Val Phe Ser Pro Lys Leu Ser

35					40					45									
Arg	Thr	Arg	Ser	Leu	Ser	Gln	Lys	Ala	Leu	Gly	Thr	Trp	Ser	Thr	Asp				
50					55					60									
Ser	Trp	Thr	Gln	Val	Ser	Leu	Pro	Cys	Arg	Gly	Ser	Pro	Ser	Ser	Glu				
65					70					75					80				
Lys	Asn	Ile	Ser	Leu	Gly	Lys	Lys	Lys	Ser	Ser	Trp	Arg	Thr	Leu	Phe				
85					90					95									
Arg	Val	Ala	Glu	Lys	Glu	Glu	Gly	Leu	Pro	Xaa	Ser	Pro	Lys	Glu	Ile				
100					105					110									
Arg	Ala	Gln	Gly	Pro	Gln	Gly	Pro	Phe	Pro	Val	Glu	Arg	Gln	Ser	Gly				
115					120					125									
Phe	His	Asn	Gln	His	Trp	Pro	Arg	Ser	Leu	Ser	Ser	Val	Glu	Gln	Pro				
130					135					140									
Trp	Arg	Val	Lys	Leu	Trp	Ile	Pro	Lys	Trp	Leu	Val	Leu	Pro	Thr	Glu				
145					150					155					160				
Trp	Leu	Lys	Leu	Phe	Thr	Pro	Gly	His	His	Gln	Met	Ser	Ser	Thr	Ala				
165					170					175									
Arg	Glu	Glu	Ala	Ser	Ser	Lys	Arg	Gly	Ser	Arg	Arg	Phe	Leu	Tyr	Phe				
180					185					190									
Arg	Phe	Glu	Gly	Pro	Trp	Asp	Ser	Lys	Asn	Lys	Asp	Gly	Glu	Asp	Gln				
195					200					205									
Ile	Ile	Ser	Lys	Ile	Val	Glu	Leu	Leu	Lys	Ser	Ser	Gly	Asp	Gln	Leu				
210					215					220									
Gly	Arg	Glu	Ile	Lys	Lys	Asp	Lys	Ala	Leu	Met	Ser	Ser	Phe	Gln	Asp				
225					230					235					240				
Gly	Leu	Ser	Tyr	Ser	Thr	Phe	Lys	Thr	Ile	Thr	Asp	Leu	Phe	Leu	Arg				
245					250					255									
Asp	Val	Asp	Thr	Arg	Gly	Glu	Ser	Glu	Val	Lys	Ala	Arg	Gly	Phe	Lys				
260					265					270									
Ala	Ala	Leu	Ala	Ile	Asp	Ala	Ile	Ala	Lys	Leu	Thr	Ala	Ser	Asp	Asn				
275					280					285									
His	Pro	Met	Asn	Arg	Met	Leu	Gly	Phe	Gly	Thr	Lys	Tyr	Leu	Lys	Glu				
290					295					300									
Tyr	Phe	Ser	Pro	Trp	Val	Gln	Gln	Asn	Gly	Gly	Trp	Glu	Lys	Ile	Leu				
305					310					315					320				
Gly	Ile	Ser	His	Glu	Glu	Val	Asp												
325																			

<210> 43
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 43
gggctcgaga tgtgcagcac cagtgtgtat g

31

<210> 44
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: synthetic
primer

<400> 44
ccaagcttta agtctacttc ttcattgtgat atccc

35

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44282 A3

(51) International Patent Classification⁷: **C12N 15/12**,
15/11, 15/62, C07K 14/47, 16/18, A01K 67/027, A61K
38/17, 48/00, C12Q 1/68, G01N 33/68, 33/53

(74) Agents: **CADENA, Deborah, L.**, et al.; Campbell & Flores
LLP, 7th floor, 4370 La Jolla Village Drive, San Diego, CA
92122 (US).

(21) International Application Number: **PCT/US00/33793**

(22) International Filing Date:
13 December 2000 (13.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/461,641 14 December 1999 (14.12.1999) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 09/461,641 (CIP)
Filed on 14 December 1999 (14.12.1999)

(71) Applicant (for all designated States except US): **THE
BURNHAM INSTITUTE** [US/US]; 10901 N. Torrey
Pines Road, La Jolla, CA 92037 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **REED, John, C.**
[US/US]; 17044 El Camino Real, P.O. Box 137, Rancho
Santa Fe, CA 92067 (US). **GODZIK, Adam** [PL/US];
9184 Buckwheat Street, San Diego, CA 92129 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
CZ (utility model), DE, DE (utility model), DK, DK (utility
model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility
model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO,
RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
21 February 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 01/44282 A3

(54) Title: **BCL-G POLYPEPTIDES, ENCODING NUCLEIC ACIDS AND METHODS OF USE**

(57) Abstract: The invention provides Bcl-G polypeptides and encoding nucleic acids. Bcl-G polypeptides include Bcl-G₁ and Bcl-G₅. The invention also provides mouse Bcl-G. The invention also provides vectors containing Bcl-G nucleic acids, host cells containing such vectors, Bcl-G anti-sense nucleic acids and related compositions. The invention additionally provides Bcl-G oligonucleotides that can be used to hybridize to or amplify a Bcl-G nucleic acid. Anti-Bcl-G specific antibodies are also provided. Further provided are kits containing Bcl-G nucleic acids or Bcl-G specific antibodies. Such kits and reagents can be used to diagnose cancer, monitor response to therapy, or predict the prognosis of a cancer patient. The invention additionally provides methods of modulating apoptosis using Bcl-G polypeptides, encoding nucleic acids, or compounds that modulate the activity or expression of Bcl-G polypeptides. The methods for modulating apoptosis can be used to treat diseases such as cancer.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/33793

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/11 C12N15/62 C07K14/47 C07K16/18
A01K67/027 A61K38/17 A61K48/00 C12Q1/68 G01N33/68
G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EMBASE, EPO-Internal, EMBL, PASCAL, MEDLINE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! 31 July 1997 (1997-07-31) MARRA M. ET AL.: " Knowles Solter mouse 2 cell Mus musculus cDNA clone " retrieved from EBI Database accession no. AA536718 XP002174348	1-6
Y	abstract	7-34, 43, 45, 46, 48, 50-53
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

10 August 2001

Date of mailing of the international search report

24/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

Int :ional Application No

PCT/US 00/33793

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! 26 April 1999 (1999-04-26) MARRA M. ET AL. : "Stratagene mouse testis (#937308) Mus musculus cDNA clone" retrieved from EBI Database accession no. AI614194 XP002174349	1-6
Y	abstract	7-34, 43, 45, 46, 48, 50-53
Y	---- WO 95 28497 A (JOLLA CANCER RES FOUND) 26 October 1995 (1995-10-26) claims 1-14; figures 5, SEQ.ID.24	7-34, 43, 45, 46, 48, 50-53
Y	---- WO 95 00642 A (ARCH DEV CORP ; NUNEZ GABRIEL (US); BOISE LAWRENCE H (US); REGENT O) 5 January 1995 (1995-01-05) abstract; claims 1-45; figures 1-4	7-34, 43, 45, 46, 48, 50-53
A	---- WO 97 45128 A (APOPTOSIS TECHNOLOGY INC) 4 December 1997 (1997-12-04) abstract; claim 2; figures 2, 8	48
A	---- US 5 691 179 A (KORSMEYER STANLEY J) 25 November 1997 (1997-11-25) column 2, paragraph 3 column 28, paragraph 3 - paragraph 4	48
T	---- GUO, BIN ET AL: "Bcl - G, a novel pro-apoptotic member of the Bcl-2 family" J. BIOL. CHEM. (2001), 276(4), 2780-2785 , XP002174347 the whole document -----	1-34, 43, 45, 46, 48, 50-53

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 35,43,44 partly and 36-40,47,49

Claims 35,43,44,46 partly and 36-40,47,49 relating to modulators to the BCL-g polypeptide could not be searched as its subject-matter was insufficiently disclosed

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/33793

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9528497 A	26-10-1995	AU 2290195 A EP 0759090 A US 5702897 A	10-11-1995 26-02-1997 30-12-1997
WO 9500642 A	05-01-1995	US 5646008 A AU 7177394 A CA 2165969 A EP 0706568 A JP 9502603 T US 5834309 A	08-07-1997 17-01-1995 05-01-1995 17-04-1996 18-03-1997 10-11-1998
WO 9745128 A	04-12-1997	US 5834234 A EP 0910387 A JP 2000510686 T US 6057132 A	10-11-1998 28-04-1999 22-08-2000 02-05-2000
US 5691179 A	25-11-1997	AU 688368 B AU 7604994 A BR 9407583 A CA 2170143 A EP 0722275 A JP 9502088 T JP 2000336100 A NZ 271929 A WO 9505750 A US 5700638 A US 5856171 A US 5834209 A US 5955595 A US 5942490 A US 6184202 B	12-03-1998 21-03-1995 07-01-1997 02-03-1995 24-07-1996 04-03-1997 05-12-2000 25-03-1998 02-03-1995 23-12-1997 05-01-1999 10-11-1998 21-09-1999 24-08-1999 06-02-2001

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.